

The Integrated Square-Root Process

by

Daniel Dufresne
University of Montreal

RESEARCH PAPER NUMBER 90

November 2001

Centre for Actuarial Studies
Department of Economics
The University of Melbourne
Victoria 3010
Australia

THE INTEGRATED SQUARE-ROOT PROCESS

Daniel Dufresne, University of Montreal

Abstract

The square-root process has been used to model interest rates and stochastic volatility. This paper studies some of its properties, particularly those of the integral of the process over time. After summarizing the properties of the square-root process, the Laplace transform of the integral of the square-root process is derived. Three methods for the computation of the moments of this integral are given, as well as some properties of the density of the integral. The last section studies the relationship between the Laplace transforms of a variable and of its reciprocal, a topic which arises in the previous analysis and elsewhere. An application to the generalized inverse Gaussian distribution is given.

Keywords: SQUARE-ROOT PROCESS; STOCHASTIC VOLATILITY; VOLATILITY SWAPS; ASIAN OPTIONS; INTEREST RATE MODELS; GENERALIZED INVERSE GAUSSIAN DISTRIBUTION

1. Introduction

The *square-root process* is the unique strong solution of the stochastic differential equation (SDE in the sequel)

$$dX = (\alpha X + \beta) dt + \gamma \sqrt{X} dW, \quad X_0 = \bar{x} \geq 0, \quad (1.1)$$

where $\alpha \in \mathbb{R}$, $\beta > 0$, $\gamma > 0$, $\bar{x} \geq 0$, and W is standard Brownian motion. Other names used for the same process are *squared Gaussian* (since the square of some Gaussian processes can be shown to satisfy (1.1) — a comment attributed to L.C.G. Rogers), *Feller* (because of the early work of William Feller on the process), *CIR* (since Cox, Ingersoll & Ross (1985) used (1.1) as a model for the spot interest rate, and gave the formulas for risk-free bonds under this assumption), and *Heston model* (since Heston (1993) studied the pricing of options when the squared volatility of the underlying security satisfies (1.1)).

The integral of X arises in both volatility and interest rate models. If the stock price follows the process

$$dS = \mu S dt + \sqrt{X} S dV,$$

where (V, W) is two-dimensional Brownian motion (possibly correlated), and X is the solution of (1.1), then the integral

$$Y_t = \int_0^t X_s ds$$

This version: 16-11-2001

is the cumulative variance of the stock returns up to time t . This integrated variance process occurs in volatility swaps (Carr, 2001), and one can imagine other derivatives based on Y_t .

The process Y_t obviously arises in bond pricing, when spot interest rates follow the CIR model. This is because the price of a risk-free zero-coupon bond maturing at time t equals

$$E_Q e^{-Y_t}$$

where Q is a risk-neutral measure. Another example would be options based on the average spot interest rate, for instance a call on the average of spot rates X_t over $[0, T]$, with strike R_0 :

$$\left(\frac{1}{T} \int_0^T X_t dt - R_0 \right)_+ = e^{-Y_t} \frac{1}{T} (Y_T - TR_0)_+$$

The price of such an option would be the expectation of the payoff with respect to the risk neutral measure, which requires knowledge of the distribution of Y_T .

The moments and distribution of Y_t may also be of interest in the statistical estimation of the parameters α, β, γ in (1.1).

Here is a summary of the paper. Section 2 shows that the moments of X_t are solutions of a sequence of ordinary differential equations, and that these moments allow the determination of the moment generating function and of the density of the process. The moments of X_t are required in the sequel, and the technique used to obtain them is used again in Section 3. Moreover, Section 2 yields an improvement of a result due to Dufresne (1989), regarding the moments of a process related to Asian options and the Courtadon interest rate model. The density of X_t in (1.1) is of course well known, but the derivation given in Section 2 is not common (the author has not seen it in the literature), and clearly shows that the distribution of the process is the convolution of two distributions, one compound Poisson/Exponential and the other gamma. (The distribution of X_t has been called "non-central chi-square," apparently because it reduces to the distribution of the square of non-zero normal variable when $\alpha = 0$.) The better known way of finding the moment generating function and density of X_t is via the relationship with Bessel processes, and this is also indicated in Section 2.

Section 3 shows how the moments of the integral of the square-root process may be obtained, either by solving linear differential equations, or more directly by a recursive procedure; the results are apparently new. Section 4 is a derivation of the Laplace transform of the integral of the square-root process, based on the formula for the price of a zero-coupon bond in the CIR framework (the latter was based on results previously known in probability theory). The Laplace transform yields a third way of calculating the moments. Section 5 extends a result which is used in Section 4, and is of independent interest in Probability Theory. Formulas are given which relate the Laplace transform of a random variable $U > 0$ to the Laplace transform of $1/U$. They extend one well-known formula for Laplace transforms. An application to the generalized inverse Gaussian distribution is given.

A subsequent paper will use the results of this one to compute the distribution of the integral of the square-root process as well as option prices.

2. A sequence of ODEs; the moments and distribution of X_t

When the square-root process (1.1) is used as a financial model, it is usually made “mean-reverting” by letting $\alpha < 0$. In the sequel no such restriction is imposed, except that Theorem 2.3 assumes $\alpha \neq 0$.

The moments of X_t are easy to obtain recursively, by applying Ito’s formula and then taking expectations. A general formula for the moments of X_t will now be derived. The technique is similar to the one used in Dufresne (1989, 1990) for the integral of geometric Brownian motion, and yields an extension of one result in Dufresne (1989) (Corollary 2.2 below). Next, the Laplace transform of the density of X_t is derived, based on the moments only. This derivation is different from the usual ones, which are either based on PDEs, or on the relationship with Bessel processes. Our derivation is based on the assumption that all the moments of X are finite. The moments are calculated and then summed to get the moment generating function (MGF). Finally, the density is exhibited, and is shown to be the convolution of a Gamma and a compound Poisson/exponential.

(N.B. The MGF of a probability measure m is the mapping

$$r \mapsto \int e^{rx} m(dx).$$

The concept is thus the same as the Laplace transform, but a + instead of a – in front of the argument.)

Applying Ito’s formula with $f(x) = x^k$ to (1.1), we get

$$dX^k = (a_k X^k + b_k X^{k-1})dt + \gamma k X^{k-\frac{1}{2}} dW_t \quad (2.1)$$

with

$$a_k = \alpha k, \quad b_k = \beta k + \frac{1}{2} \gamma^2 k(k-1).$$

Eq.(2.1) is the same as

$$X_t = \bar{x} + \int_0^t (a_k X_s^k + b_k X_s^{k-1}) ds + \int_0^t \gamma k X_s^{k-\frac{1}{2}} dW_s.$$

Here we let the initial value of the process be $\bar{x} \geq 0$, and we take as given that $E X^n < \infty$ for all $n \geq 0$. The consequence is that the Ito integral above has expected value 0. Defining $m_k(t) = E X_t^k$, $k = 0, 1, \dots$, we can then differentiate with respect to t on each side to obtain

$$m'_k(t) = a_k m_k(t) + b_k m_{k-1}(t), \quad k \geq 1, \quad t > 0 \quad (2.2)$$

$$m_k(0) = \bar{x}^k, \quad k \geq 0. \quad (2.3)$$

Eq. (2.2) is of the same type as Eq.(18) in Dufresne (1989), and can be solved in the same way. The main difference here is that the initial value of $m_k(t)$ is not 0.

Theorem 2.1. Let $a_0 = 0$ and suppose the numbers $\{a_0, \dots, a_K\}$ are distinct. Then the solution of (2.2), subject to (2.3) and $m_0(t) \equiv 1$ is

$$m_k(t) = \sum_{j=0}^k d_{kj} e^{a_j t}, \quad k = 0, \dots, K, \quad (2.4)$$

where

$$d_{kj} = \sum_{i=0}^j \bar{x}^i \left(\prod_{m=i+1}^k b_m \right) \prod_{\substack{\ell=i \\ \ell \neq j}}^k \frac{1}{a_j - a_\ell}, \quad j = 0, \dots, k. \quad (2.5)$$

Proof. Define a product over an empty set of indices as 1. Then formulas (2.4) and (2.5) are correct for $k = 0$. For $k = 1$, (2.2) implies

$$m_1(t) = m_1(0)e^{a_1 t} + b_1 e^{a_1 t} \int_0^t e^{-a_1 s} ds = \bar{x} e^{a_1 t} + b_1 \frac{e^{a_1 t} - 1}{a_1} = -\frac{b_1}{a_1} + \left(\frac{b_1}{a_1} + \bar{x} \right) e^{a_1 t}.$$

For $j = 0$ and $j = 1$, $m_j(t)$ is a combination of $e^{a_\ell t}$, $\ell \leq j$. If this claim is correct for $j = 0, \dots, k-1$, then $m_k(t)$ is the solution of an inhomogeneous ordinary differential equation with a forcing term equal to a combination of exponentials:

$$\begin{aligned} m'_k(t) - a_k m_k(t) &= \sum_{j=0}^{k-1} C_j e^{a_j t} \\ \Rightarrow m_k(t) &= m_k(0) e^{a_k t} + e^{a_k t} \int_0^t e^{-a_k s} \sum_{j=0}^{k-1} C_j e^{a_j s} ds. \end{aligned} \quad (2.6)$$

Since the $\{a_0, \dots, a_K\}$ are distinct, $m_k(t)$ has to be of the form

$$d_{k0} + d_{k1} e^{a_1 t} + \dots + d_{kk} e^{a_k t}.$$

By induction, this is true for $k = 0, \dots, K$. As a function of \bar{x} , $m_k(t)$ is a polynomial of degree k , and the coefficient of \bar{x}^k in $m_k(t)$ is $e^{a_k t}$ (because \bar{x}^k is the initial condition $m_k(0)$).

Insert (2.4) into (2.2) to obtain

$$\sum_{j=0}^k a_j d_{kj} e^{a_j t} = \sum_{j=0}^k a_k d_{kj} e^{a_j t} + \sum_{j=0}^{k-1} b_k d_{k-1,j} e^{a_j t},$$

which implies

$$d_{kj} = \frac{b_k}{a_j - a_k} d_{k-1,j} = \frac{b_k}{a_j - a_k} \dots \frac{b_{j+1}}{a_j - a_{j+1}} d_{jj}, \quad 0 \leq j < k. \quad (2.7)$$

The integrated square-root process

Since $m_j(t)$ is a polynomial of degree j in \bar{x} , the last equation implies that d_{kj} is also a polynomial of degree j in \bar{x} , which we write as

$$d_{kj} = \sum_{i=0}^j d_{kji} \bar{x}^i.$$

By identifying the coefficients of \bar{x}^i in (2.7), we get

$$d_{kji} = \frac{b_k}{a_j - a_k} \cdots \frac{b_{j+1}}{a_j - a_{j+1}} d_{jji}, \quad 0 \leq i \leq j < k.$$

We also know that $d_{kkk} = 1$, since the leading coefficient of $m_k(t)$, as a polynomial in \bar{x} , is $e^{a_k t}$ (see (2.6)). The missing constants are thus d_{kki} , $i = 0, \dots, k-1$, $k \geq 1$. All this shows that the problem is the same for each power of \bar{x} : for any $i = 0, 1, \dots$, we have

$$d_{iii} = 1 \tag{2.8a}$$

$$d_{kji} = \frac{b_k}{a_j - a_k} d_{k-1,j,i}, \quad i \leq j < k \tag{2.8b}$$

$$\sum_{j=i}^k d_{kji} = 0, \quad 0 \leq i < k. \tag{2.8c}$$

(The last equality comes from the initial condition $m_k(0) = \bar{x}^k$.) Hence, the problem of finding $\{d_{kj0}; 0 \leq j \leq k\}$ when $\bar{x} \neq 0$ is the same as finding $\{d_{kj}; 0 \leq j \leq k\}$ when $\bar{x} = 0$. The latter was done in Dufresne (1989), for the special case $b_k = k$. The same arguments work, however, for a general sequence $\{b_k; k \geq 1\}$. The main argument is that

$$\sum_{j=0}^k \prod_{\substack{\ell=0 \\ \ell \neq j}}^k \frac{1}{a_j - a_\ell} = 0 \tag{2.9}$$

for any distinct numbers $\alpha_0, \dots, \alpha_k$. We recall the proof given in Dufresne (1989). Lagrange's formula for partial fractions decomposition of a rational function is

$$\frac{P(x)}{Q(x)} = \sum_{j=0}^k \frac{1}{x - a_j} \frac{P(a_j)}{Q'(a_j)},$$

for polynomials P and Q , the latter with $k+1$ distinct zeros $\{a_0, \dots, a_k\}$, the former of degree less than or equal to k . Let

$$Q(x) = \prod_{j=0}^k (x - a_j), \quad P(x) = \frac{y - x}{Q(y)}.$$

Then

$$P(x) = \sum_{j=0}^k \frac{y - a_j}{Q(y)} \frac{Q(x)}{(x - a_j)Q'(a_j)}.$$

Letting $y \rightarrow x$ yields (2.9).

For $i = j = 0$, we find (from (2.8a, b))

$$d_{k00} = \left(\prod_{m=1}^k b_m \right) \prod_{\substack{\ell=0 \\ \ell \neq 0}}^k \frac{1}{a_0 - a_\ell}, \quad k \geq 0.$$

Next, $d_{110} = -d_{100} = b_1/(a_1 - a_0)$ (from (2.8c)), which implies

$$d_{k10} = \frac{b_k \cdots b_2}{(a_1 - a_k) \cdots (a_1 - a_2)} d_{110} = \left(\prod_{m=1}^k b_m \right) \prod_{\substack{\ell=0 \\ \ell \neq 1}}^k \frac{1}{a_1 - a_\ell}, \quad k \geq 1.$$

To prove the result in general, represent the $\{d_{kj0}\}$ on a grid, with k horizontal and j vertical. For some integer $J > 0$, suppose that

$$d_{kj0} = \left(\prod_{m=1}^k b_m \right) \prod_{\substack{\ell=0 \\ \ell \neq j}}^k \frac{1}{a_j - a_\ell}, \quad k \geq j, \quad j = 0, \dots, J-1, \quad (2.10)$$

that is, assume the result holds for the lines $j = 0, \dots, J-1$. Then, by (2.8c) and (2.9), (2.10) also holds for $k = j = J$. From this and (2.8b), we find, for $k > J$,

$$d_{kJ0} = \frac{b_k \cdots b_{J+1}}{(a_J - a_k) \cdots (a_J - a_{J+1})} d_{JJ0} = \left(\prod_{m=1}^k b_m \right) \prod_{\substack{\ell=0 \\ \ell \neq J}}^k \frac{1}{a_J - a_\ell}.$$

By induction, (2.10) applies for all $0 \leq j \leq k$. For the coefficients of \bar{x}^i , $i \geq 1$, the problem is the same, except that it “starts” on line i , so that, in the above arguments, (j, k) has to be replaced with $(j + i, k + i)$. \square

Remark. Similar formulas hold when some of the $\{a_0, \dots, a_K\}$ are identical. The only difference is that terms of the form $t^\ell e^{a_j t}$ make their appearance, where ℓ is an integer which depends on the number of times the same a_k appears in $\{a_0, \dots, a_K\}$. For instance, suppose $a_0 = 0, a_1 = a_2 \neq 0$. The solution of (2.2) is then

$$m_2(t) = \bar{x}^2 e^{a_1 t} + b_2 \left(\bar{x} + \frac{b_1}{a_1} \right) t e^{a_1 t} - \frac{b_1 b_2}{a_1^2} (e^{a_1 t} - 1).$$

The $t e^{a_1 t}$ reappears in $m_j(t)$, $j \geq 3$. The solutions of the differential equations nevertheless depend continuously on the parameters a_0, a_1, \dots , so that the solutions when some of them are identical can be found by taking limits in the solutions given in the theorem. \square

The integrated square-root process

As a first application of Theorem 2.1, we generalize Proposition 2 of Dufresne (1989). Suppose $\{S_t\}$ satisfies

$$dS_t = (\eta S_t + \zeta) dt + \sigma S_t dW_t, \quad S_0 = \bar{x}. \quad (2.11)$$

Then the moments $m_k(t) = \mathbb{E} S_t^k$, $k = 0, 1, \dots$, satisfy Eqs.(2.3-2.4), with

$$a_k = k\eta + k(k-1)\sigma^2/2, \quad b_k = k\zeta. \quad (2.12)$$

Corollary 2.2. *If the $\{a_k\}$ are all distinct, the moments of the process in (2.11) are given by (2.4), with $\{a_k\}$ as in (2.12) and*

$$d_{kj} = k! \sum_{i=0}^j \frac{\bar{x}^i \zeta^{k-i}}{i!} \prod_{\substack{\ell=i \\ \ell \neq j}}^k \frac{1}{a_j - a_\ell}, \quad j = 0, \dots, k.$$

Remark. The process S in (2.11) has been called the Courtadon interest rate model (Courtadon, 1982). The moments of the spot rate are thus given by Corollary 2.2. The limit distribution of S_t as t tends to infinity (inverse Gamma if $\eta < \sigma^2/2$, ∞ otherwise) is derived in Dufresne (1990). In that paper, S_t represents the market value of a an initial amount S_0 and of a continuous payment stream of ζ units, all invested in a security represented by a geometric Brownian motion. \square

Theorem 2.3. *Suppose $\alpha \neq 0$. If X_t satisfies (1.1), then its moments are*

$$\mathbb{E} X_t^k = \sum_{j=0}^k \theta_{kj} e^{\alpha j t}, \quad k = 0, 1, \dots,$$

where

$$\theta_{kj} = \sum_{i=0}^j \bar{x}^i \frac{k! (-1)^{k-j} \bar{u}^{k-i} (\bar{v})_k}{i! (j-i)! (k-j)! (\bar{v})_i}, \quad 0 \leq j \leq k$$

$$\bar{u} = \frac{\gamma^2}{2\alpha}, \quad \bar{v} = \frac{2\beta}{\gamma^2}$$

$$(y)_0 = 1, \quad (y)_k = y(y+1)\cdots(y+k-1), \quad k \geq 1.$$

Proof. We have

$$\begin{aligned} \prod_{m=1}^k b_m &= k! \left[\frac{\gamma^2}{2} k + \left(\beta - \frac{\gamma^2}{2} \right) \right] \cdots \left[\frac{\gamma^2}{2} 1 + \left(\beta - \frac{\gamma^2}{2} \right) \right] \\ &= k! \left(\frac{\gamma^2}{2} \right)^k \left[\frac{2\beta}{\gamma^2} + k - 1 \right] \cdots \frac{2\beta}{\gamma^2} \\ &= k! \left(\frac{\gamma^2}{2} \right)^k (\bar{v})_k. \end{aligned}$$

For $i \leq j$,

$$\begin{aligned} \prod_{\substack{\ell=i \\ \ell \neq j}}^k (a_j - a_\ell) &= \alpha^{k-i} \prod_{\ell=i}^{j-1} (j - \ell) \prod_{\ell=j+1}^k (j - \ell) \\ &= (-1)^{k-j} \alpha^{k-i} (j - i)! (k - j)!. \end{aligned}$$

Hence

$$\left(\prod_{m=i+1}^k b_m \right) \prod_{\substack{\ell=i \\ \ell \neq j}}^k \frac{1}{a_j - a_\ell} = \frac{k!}{i!} \left(\frac{\gamma^2}{2\alpha} \right)^{k-i} \frac{(\bar{v})_k}{(\bar{v})_i} \frac{(-1)^{k-j}}{(j - i)! (k - j)!}. \quad \square$$

The MGF of X_t will now be obtained by summing the moments of X_t . We begin by assuming that the following sum converges:

$$\sum_{k=0}^{\infty} \frac{s^k}{k!} \mathbb{E} X_t^k = \sum_{j=0}^{\infty} e^{a_j t} \sum_{k=j}^{\infty} \frac{s^k}{k!} \theta_{kj} = \sum_{k=0}^{\infty} \frac{s^k}{k!} \sum_{j=0}^k e^{a_j t} \sum_{i=0}^j \bar{x}^i \frac{k! (-1)^{k-j} \bar{u}^{k-i}}{i! (j - i)! (k - j)!} \frac{(\bar{v})_k}{(\bar{v})_i}. \quad (2.13)$$

This sum will be evaluated by summing first over k , then over j , and finally over i . This procedure needs to be justified, as the summands are not all of the same sign. The justification is given below. We will use the formula

$$(1 - y)^{-c} = \sum_{n=0}^{\infty} (c)_n \frac{y^n}{n!}, \quad (2.14)$$

which is valid for $c \in \mathbb{R}$, $|y| < 1$.

For fixed $i \leq j$ and small enough $|s|$, we have

$$\sum_{k=j}^{\infty} \bar{x}^i \frac{e^{a_j t} s^k}{k!} \frac{k! (-1)^{k-j} \bar{u}^{k-i}}{i! (j - i)! (k - j)!} \frac{(\bar{v})_k}{(\bar{v})_i} = \bar{x}^i \frac{(\bar{v})_j}{(\bar{v})_i} \frac{s^j \bar{u}^{j-i} e^{a_j t}}{i! (j - i)!} (1 + s\bar{u})^{-\bar{v}-j}.$$

Next,

$$\begin{aligned} \bar{x}^i \sum_{j=i}^{\infty} \frac{(\bar{v})_j}{(\bar{v})_i} \frac{s^j \bar{u}^{j-i} e^{a_j t}}{i! (j - i)!} (1 + s\bar{u})^{-\bar{v}-j} &= \frac{s^i \bar{x}^i e^{\alpha i t}}{i!} (1 + s\bar{u})^{-\bar{v}-i} \sum_{j=i}^{\infty} \frac{(\bar{v})_j}{(\bar{v})_i} \frac{s^{j-i} \bar{u}^{j-i} e^{\alpha t (j-i)}}{(j - i)!} (1 + s\bar{u})^{i-j} \\ &= \frac{s^i \bar{x}^i e^{\alpha i t}}{i!} (1 + s\bar{u})^{-\bar{v}-i} \left(1 - \frac{s\bar{u} e^{\alpha t}}{1 + s\bar{u}} \right)^{-\bar{v}-i} \\ &= \frac{s^i \bar{x}^i e^{\alpha i t}}{i!} [1 - s\bar{u}(e^{\alpha t} - 1)]^{-\bar{v}-i}. \end{aligned}$$

Finally, summing over i ,

$$\mathbb{E} e^{sX_t} = [1 - s\bar{u}(e^{\alpha t} - 1)]^{-\bar{v}} \exp \left[\frac{s\bar{x}e^{\alpha t}}{1 - s\bar{u}(e^{\alpha t} - 1)} \right] = \phi(s)^{\bar{v}} \exp[\lambda_t(\phi(s) - 1)]. \quad (2.15)$$

Here

$$\phi(s) = (1 - s\mu_t)^{-1}, \quad \mu_t = \frac{\gamma^2}{2} \left(\frac{e^{\alpha t} - 1}{\alpha} \right), \quad \lambda_t = \frac{2\alpha\bar{x}}{\gamma^2(1 - e^{-\alpha t})}.$$

The function $\phi(s)$ is the MGF of an exponential distribution, with mean equal to μ_t . The distribution of X_t is thus the distribution of $X'_t + X''_t$, where the two variables are independent, $X'_t \sim \mathbf{Gamma}(\bar{v}, \mu_t)$, and X''_t is compound Poisson

$$X''_t = \sum_{i=1}^N U_i,$$

with $N \sim \mathbf{Poisson}(\lambda_t)$, $U_i \sim \mathbf{Exp}(\mu_t)$.

We now show that the sum in (2.13) converges. What we have just done is to sum the quantities

$$c_{ijk} = \frac{s^k \bar{x}^i (-1)^{k-j} \bar{u}^{k-i} e^{a_j t} (\bar{v})_k}{i!(j-i)!(k-j)! (\bar{v})_i}, \quad 0 \leq i \leq j \leq k < \infty.$$

The same steps show that

$$\sum_{0 \leq i \leq j \leq k < \infty} |c_{ijk}| = [1 - s\bar{u}(e^{\alpha t} + 1)]^{-\bar{v}} \exp \left[\frac{s\bar{x}e^{\alpha t}}{1 - s\bar{u}(e^{\alpha t} + 1)} \right],$$

if $0 \leq |s| < 1/[\bar{u}(e^{\alpha t} + 1)]$. Therefore: (1) the sum in (2.13) is convergent, (2) the order of summation of the c_{ijk} is irrelevant, and (3) the result is an analytic function of s , at least for $0 \leq |s| < 1/[\bar{u}(e^{\alpha t} + 1)]$. Expression (2.15) is therefore correct for $s < 1/[\bar{u}(e^{\alpha t} - 1)]$, by analytic continuation.

Obviously, the limit distribution of X_t as $t \rightarrow \infty$ exists if, and only if, $\alpha < 0$, in which case it is $\mathbf{Gamma}(2\beta/\gamma^2, -\gamma^2/(2\alpha))$ (the compound Poisson part, which depends on the initial condition \bar{x} , disappears in the limit). Finally, the density of X_t may be obtained by conditioning on N . To simplify the algebra, consider $U = X_t/\mu_t$: with probability $e^{-\lambda_t} \lambda_t^n/n!$, the distribution of U is $\mathbf{Gamma}(\bar{v} + n, 1)$, for $n = 0, 1, \dots$. Hence, the density of U is

$$\begin{aligned} f_U(y) &= \sum_{n=0}^{\infty} e^{-\lambda_t} \frac{\lambda_t^n}{n!} \frac{y^{\bar{v}+n-1}}{\Gamma(\bar{v}+n)} e^{-y} \mathbf{1}_{\{y>0\}} = \left(\frac{y}{\lambda_t} \right)^{\frac{\bar{v}-1}{2}} e^{-\lambda_t-y} \sum_{n=0}^{\infty} \frac{(\sqrt{\lambda_t y})^{\bar{v}-1+2n}}{n! \Gamma(\bar{v}+n)} \mathbf{1}_{\{y>0\}} \\ &= \left(\frac{y}{\lambda_t} \right)^{\frac{\bar{v}-1}{2}} e^{-\lambda_t-y} I_{\bar{v}-1}(2\sqrt{\lambda_t y}) \mathbf{1}_{\{y>0\}}, \end{aligned}$$

where I_ν is the modified Bessel function of the first kind of argument ν (Lebedev, 1972, p.108):

$$I_\nu(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{\nu+2k}}{\Gamma(k+1)\Gamma(k+\nu+1)}, \quad |\arg z| < \pi, \quad \nu \in \mathbb{C}. \quad (2.16)$$

The density of X_t is thus

$$f_{X_t}(x) = \frac{1}{\mu_t} \left(\frac{x e^{-\alpha t}}{\bar{x}} \right)^{\frac{\bar{\nu}-1}{2}} e^{-\lambda_t - x/\mu_t} I_{\bar{\nu}-1} \left(\frac{4\alpha}{\gamma^2(e^{\alpha t} - 1)} \sqrt{x \bar{x} e^{\alpha t}} \right) \mathbf{1}_{\{x>0\}}. \quad (2.17)$$

(When $\bar{x} = 0$ this reduces to the **Gamma**($\bar{\nu}, \mu_t$) density.)

The MGF and the density above may also be obtained from the relationship between the solution of the SDE (1.1) and Bessel processes (see Revuz & Yor, 1999, Chapter 11). For a constant $p > 0$, the process $\tilde{X}_t = pX_t$ satisfies

$$\tilde{X}_t = p\bar{x} + \int_0^t (\alpha \tilde{X}_s + \beta p) ds + \gamma \sqrt{p} \int_0^t \sqrt{\tilde{X}_s} dW_s$$

Letting $p = 4/\gamma^2$, we find that

$$\tilde{X}_t = \tilde{x} + \int_0^t (\alpha \tilde{X}_s + \tilde{\beta}) ds + 2 \int_0^t \sqrt{\tilde{X}_s} dW_s,$$

where $\tilde{x} = 4\bar{x}/\gamma^2$, $\tilde{\beta} = 4\beta/\gamma^2$. Now, consider the square of a δ -dimensional Bessel process, that is, the unique strong solution of

$$Z_t = Z_0 + \delta t + 2 \int_0^t \sqrt{Z_s} dB_s$$

(B standard Brownian motion), and let $g(t) = (1 - e^{-\alpha t})/\alpha$ (see Revuz & Yor, 1999, Exercise (1.13), p.448). Integrating by parts, and using the time change formula for stochastic integrals (Revuz & Yor, 1999, p.180), we find

$$\begin{aligned} e^{\alpha t} Z_{g(t)} &= e^{\alpha t} \left[Z_0 + \delta g(t) + 2 \int_0^t \sqrt{Z_{g(s)}} dB_{g(s)} \right] \\ &= Z_0 + \alpha \int_0^t e^{\alpha s} Z_{g(s)} ds + \delta \int_0^t e^{\alpha s} dg(s) + 2 \int_0^t \sqrt{e^{\alpha s} Z_{g(s)}} e^{\alpha s/2} dB_{g(s)}. \end{aligned}$$

The process

$$\tilde{B}_t = \int_0^t e^{\alpha s/2} dB_{g(s)}$$

is a continuous martingale with quadratic variation equal to t , and is thus standard Brownian motion (with respect to its own filtration). Hence, the process $\tilde{Z}_t = e^{\alpha t} Z_{g(t)}$ satisfies

$$\tilde{Z}_t = Z_0 + \int_0^t (\alpha \tilde{Z}_s + \delta) dt + 2 \int_0^t \sqrt{\tilde{Z}_s} d\tilde{B}_s.$$

Consequently, the processes $\{\tilde{Z}_t; t \geq 0\}$ and $\{\tilde{X}_t; t \geq 0\}$ have the same distribution, if

$$Z_0 = \tilde{x} = \frac{4\bar{x}}{\gamma^2}, \quad \delta = \tilde{\beta} = \frac{4\beta}{\gamma^2}.$$

Finally, the process $\{X_t; t \geq 0\}$ has the same distribution as $\{\frac{\gamma^2}{4}e^{\alpha t}Z_{g(t)}; t \geq 0\}$. The MGF of Z_τ is (Revuz & Yor, 1999, p.441)

$$\mathbb{E}e^{\rho Z_\tau} = (1 - 2\rho\tau)^{-\delta/2} \exp[\rho Z_0/(1 - 2\rho\tau)].$$

It can be checked that appropriate substitutions yield (2.15). In the same fashion, the density of Z_τ is

$$f_{Z_\tau}(z) = \frac{1}{2\tau} \left(\frac{z}{Z_0}\right)^{\frac{1}{2}(\frac{\delta}{2}-1)} e^{-(Z_0+z)/2\tau} I_{\frac{\delta}{2}-1} \left(\frac{\sqrt{Z_0 z}}{\tau}\right) \mathbf{1}_{\{z>0\}}.$$

This implies that the density of X_t is (for $x > 0$)

$$\begin{aligned} & \frac{2\alpha e^{-\alpha t}}{\gamma^2(1 - e^{-\alpha t})} \left(\frac{x e^{-\alpha t}}{\bar{x}}\right)^{\frac{1}{2}(\bar{\nu}-1)} \exp\left[-\left(\frac{4\bar{x}}{\gamma^2} + \frac{4x e^{-\alpha t}}{\gamma^2}\right)\alpha/[2(1 - e^{-\alpha t})]\right] I_{\bar{\nu}-1} \left(\frac{4\alpha\sqrt{x\bar{x}e^{-\alpha t}}}{\gamma^2(1 - e^{-\alpha t})}\right) \\ &= \frac{1}{\mu_t} \left(\frac{x e^{-\alpha t}}{\bar{x}}\right)^{\frac{1}{2}(\bar{\nu}-1)} e^{-\lambda_t - x/\mu_t} I_{\bar{\nu}-1} \left(\frac{4\alpha\sqrt{x\bar{x}e^{-\alpha t}}}{\gamma^2(1 - e^{-\alpha t})}\right), \end{aligned}$$

which agrees with (2.17).

It is also possible to find an expression for $\mathbb{E}X_t^p$ for non-integral p . One could integrate the density times x^p term by term; an alternative is the following: since

$$\frac{1}{y^q} = \frac{1}{\Gamma(q)} \int_0^\infty s^{q-1} e^{-sy} ds, \quad q, y > 0, \quad (2.18)$$

we have

$$\begin{aligned} \mathbb{E}X_t^{-q} &= \frac{1}{\Gamma(q)} \int_0^\infty s^{q-1} \mathbb{E}e^{-sX_t} ds \\ &= \frac{1}{\Gamma(q)} \int_0^\infty s^{q-1} \phi(-s)^{\bar{\nu}} e^{\lambda_t(\phi(-s)-1)} ds \\ &= \frac{\mu_t^{-q} e^{-\lambda_t}}{\Gamma(q)} \int_0^1 r^{\bar{\nu}-q-1} (1-r)^{q-1} e^{\lambda_t r} dr \end{aligned} \quad (2.19)$$

$$= \mu_t^{-q} e^{-\lambda_t} \frac{\Gamma(\bar{\nu}-q)}{\Gamma(\bar{\nu})} {}_1F_1(\bar{\nu}-q, \bar{\nu}; \lambda_t), \quad (2.20)$$

where ${}_1F_1$ is the confluent hypergeometric function

$${}_1F_1(r, s; z) = \sum_{n=0}^{\infty} \frac{(r)_n z^n}{(s)_n n!}, \quad r, s, z \in \mathbb{C}, \quad s \neq 0, -1, \dots$$

Here we have used the integral representation (9.11.1), p.266 of Lebedev (1972) for the confluent hypergeometric function. The above calculation is seen to be correct for $0 \leq q < \bar{v}$, while $\mathbb{E}X_t^{-q}$ is infinite for $q \leq \bar{v}$ (this is because the integral in (2.19) tends to infinity as $q \uparrow \bar{v}$). Since $\mathbb{E}X_t^{-q}$ is an analytic function of q , and (2.20) is analytic for $\text{Re}(-q) > -\bar{v}$, we get the following result.

Theorem 2.4. $\mathbb{E}X_t^p = \mu_t^p e^{-\lambda t} \frac{\Gamma(\bar{v} + p)}{\Gamma(\bar{v})} {}_1F_1(\bar{v} + p, \bar{v}; \lambda t), \quad p > -2\beta/\gamma^2.$

This formula may be reconciled with the one in Theorem 2.3 by appealing to the identity (Lebedev, 1972, p.267)

$${}_1F_1(r, s; z) = e^z {}_1F_1(s - r, s; -z), \quad s \neq 0, -1, \dots$$

If $p = k$, a non-negative integer, then $\mathbb{E}X_t^k$ may be expressed as a series which terminates:

$$\begin{aligned} \mathbb{E}X_t^k &= \mu_t^k \frac{\Gamma(\bar{v} + k)}{\Gamma(\bar{v})} {}_1F_1(-k, \bar{v}; -\lambda t) \\ &= \sum_{i=0}^k \frac{(-1)^i (-k)_i (\bar{v})_k \mu_t^i \lambda_t^i}{(\bar{v})_i i!} \mu_t^{k-i} \\ &= \sum_{i=0}^k \frac{k! (\bar{v})_k}{(k-i)! (\bar{v})_i} \frac{\bar{x}^i e^{\alpha i t}}{i!} \bar{u}^{k-i} \sum_{\ell=0}^{k-i} \binom{k-i}{\ell} e^{\alpha \ell t} (-1)^{k-i-\ell} \\ &= \sum_{i=0}^k \frac{k! (\bar{v})_k}{(k-i)! (\bar{v})_i} \frac{\bar{x}^i e^{\alpha i t}}{i!} \bar{u}^{k-i} \sum_{j=i}^k \frac{(k-i)!}{(j-i)! (k-j)!} e^{\alpha(j-i)t} (-1)^{k-j} \\ &= \sum_{j=0}^k e^{\alpha j t} \sum_{i=0}^j \bar{x}^i \frac{k! (-1)^{k-j} \bar{u}^{k-i} (\bar{v})_k}{i! (j-i)! (k-j)! (\bar{v})_i}. \end{aligned}$$

3. Moments of the integrated square-root process

In this section, two techniques are given for the calculation of the moments of the integral of the square-root process $\{Y_t\}$.

Suppose X is the solution of SDE (1.1), and define

$$Y_t = \int_0^t X_s ds, \quad M_{jk}(t) = \mathbb{E}Y_t^j X_t^k,$$

for non-negative integers j, k ; M_{jk} is finite for all $t \geq 0$, since Y_t has finite moments of all order (since the Laplace transform of Y_t exists in a neighbourhood of the origin, see Section 4). Applying Ito's formula, we get

$$\begin{aligned} d(Y_t^j X_t^k) &= X_t^k d(Y_t^j) + Y_t^j d(X_t^k) \\ &= jY_t^{j-1} X_t^{k+1} dt + Y_t^j (a_k X_t^k + b_k X_t^{k-1}) dt + \gamma k Y_t^j X_t^{k-\frac{1}{2}} dW_t. \end{aligned}$$

Since $\mathbb{E}X_t^p < \infty$ for all positive p , it follows that

$$\mathbb{E} \int_0^t \left(\gamma k Y_s^j X_s^{k-\frac{1}{2}} \right)^2 ds < \infty$$

for all $j, k \in \mathbb{N}$, and so

$$\mathbb{E} \int_0^t \gamma k Y_s^j X_s^{k-\frac{1}{2}} dW_s = 0.$$

This implies

$$\frac{d}{dt} M_{jk}(t) = a_k M_{jk}(t) + b_k M_{j,k-1}(t) + j M_{j-1,k+1}(t). \quad (3.1)$$

These are linear ordinary differential equations with constant coefficients, and their solutions are straightforward (though tedious to obtain by hand). In order to find $M_{j0}(t)$, it is necessary to calculate $M_{j-1,1}(t)$, $M_{j-2,2}(t)$, and so on. If we represent the $\{M_{jk}\}$ on a grid, with j horizontal and k vertical, M_{j0} is obtained after solving the differential equations on the diagonals $M_{0,i}, \dots, M_{i,0}$ for $i \leq j$. The following result is proved by suitably adapting the arguments used in Section 2.

Theorem 3.1. *Suppose $\alpha \neq 0$, and that X satisfies (1.1). Then, for $j, k \in \mathbb{N}$,*

$$M_{jk}(t) = \sum_{m=0}^{j+k} M_{jkm}(t) e^{\alpha m t}, \quad (3.2)$$

where $M_{jkm}(t)$ is a polynomial in t with degree

$$\deg M_{jkm} \leq \begin{cases} j & \text{if } 0 \leq m \leq k \\ j+k-m & \text{if } k+1 \leq m \leq j+k. \end{cases} \quad (3.3)$$

Proof. We proceed by induction on j and then on k . The theorem is correct for $j = 0$ and all $k \geq 0$ by Theorem 2.3. Suppose (3.2)-(3.3) are correct for $j = 0, \dots, J-1$ and all $k \geq 0$, for some $J \geq 1$. Based on this assumption, the solution of (3.1) is

$$M_{Jk} = e^{a_k t} M_{Jk}(0) + e^{a_k t} \int_0^t e^{-a_k s} R_{Jk}(s) ds, \quad (3.4)$$

where

$$\begin{aligned} R_{Jk}(t) &= b_k M_{J,k-1}(t) + J M_{J-1,k+1}(t) \\ &= \sum_{m=0}^{J+k-1} [b_k M_{J,k-1,m}(t) + J M_{J-1,k+1,m}(t)] e^{a_m t} + J M_{J-1,k+1,J+k}(t) e^{a_{J+k} t}. \end{aligned}$$

We show that this implies that (3.2)-(3.3) hold for $j = J, k = 0$. Since $a_0 = b_0 = M_{J0}(0) = 0$, (3.4) reduces to

$$M_{J0}(t) = J \sum_{m=0}^J \int_0^t M_{J-1,1,m}(s) e^{a_m s} ds. \quad (3.5)$$

The integrated square-root process

Consider the terms corresponding to $m = 1, \dots, J$ in the sum above. Recall that $a_m = \alpha m \neq 0$ and that, for any constant $A \neq 0$,

$$\int_0^t e^{As} s^n ds$$

is the sum of a constant and of a polynomial of degree n times e^{At} . Hence

$$\int_0^t M_{J-1,1,m}(s) e^{a_m s} ds$$

is the sum of a constant and of a polynomial of the same degree as $M_{J-1,1,m}$ times $e^{a_m t}$.

Since $a_0 = 0$, the first term on the right hand side of (3.5) is seen to be a polynomial of degree equal to one plus the degree of $M_{J-1,1,0}$.

The above considerations imply that

$$M_{J0}(t) = \sum_{m=0}^J M_{J0m}(t) e^{\alpha m t},$$

where the functions $M_{J0m}(t)$ are polynomials in t . Moreover, the induction assumption says in particular that

$$\deg M_{J-1,1,m}(t) \leq \begin{cases} J-1 & \text{if } 0 \leq m \leq 1 \\ J-m & \text{if } 2 \leq m \leq J \end{cases}$$

and, consequently, what we have just seen also means that

$$\deg M_{J0m} \leq \begin{cases} J & \text{if } m = 0 \\ J-m & \text{if } 1 \leq m \leq J. \end{cases}$$

Keeping J fixed, make another induction assumption: (3.2)-(3.3) hold for $j = J$ and $k = 0, \dots, K-1$, for some $K \geq 1$. The proof will be finished once we show that (3.2)-(3.3) hold for $j = J$ and $k = K$ as well. By the induction assumptions and (3.4),

$$M_{JK}(t) = e^{\alpha K t} \left\{ \sum_{m=0}^{J+K-1} \int_0^t [b_m M_{J,K-1,m}(s) + J M_{J-1,K+1,m}(s)] e^{(a_m - \alpha K)s} ds + \int_0^t J M_{J-1,K+1,J+K}(s) e^{(a_{J+K} - \alpha K)s} ds \right\}. \quad (3.6)$$

By the same reasoning as before, we see that M_{JK} is of the form

$$M_{JK}(t) = \sum_{m=0}^{J+K} M_{JKm}(t) e^{\alpha m t},$$

The integrated square-root process

where $M_{JKm}(t)$ is a polynomial. In (3.6), the terms of the sum inside the curly brackets with $m \neq K$ reduce to

$$\text{constant} + \text{polynomial} \times e^{(a_m - a_K)t}$$

The degree of the polynomial is the same as that of

$$b_k M_{J,K-1,m}(s) + JM_{J-1,K+1,m}(s),$$

that is, no larger than

$$\max[\min(J, J + K - 1 - m), \min(J - 1, J + K - m)] = \min(J, J + K - m). \quad (3.7)$$

The same applies to the term following the sum, which corresponds to $m = J + K$. For the term $m = K$ inside the curly brackets, the integral reduces to a polynomial of degree no larger than

$$1 + \max[\deg(M_{J,K-1,K}), \deg(M_{J-1,K+1,K})] = 1 + \max(J - 1, J - 1) = J. \quad (3.8)$$

All these integrals are then multiplied by $e^{a_K t}$, so that (3.7) becomes the degree of M_{JKm} for $m \neq K$, and (3.8) becomes the degree of M_{JKK} . This ends the proof. \square

A general simplified formula for the polynomials M_{jkm} has not been found, but they can be calculated recursively, as the next theorem shows.

Theorem 3.2. *If we let*

$$M_{jkm}(t) = \sum_{n=0}^{j \wedge (j+k-m)} M_{jkmn} t^n,$$

$$R_{jkmn} = b_k M_{j,k-1,m,n} + j M_{j-1,k+1,m,n},$$

then

$$M_{jkmn} = - \sum_{i=n}^{j \wedge (j+k-m)} \frac{(n+1)_{i-n}}{[\alpha(k-m)]^{i-n+1}} R_{jkm i}, \quad k \neq m \quad (3.9)$$

$$M_{jkkn} = -\frac{1}{n} R_{j,k,k,n-1}, \quad n = 1, \dots, j \quad (3.10)$$

$$M_{jkk0} = - \sum_{\substack{m=0 \\ m \neq k}}^{j+k} M_{jkm0}, \quad j \geq 1 \quad (3.11)$$

$$M_{0km0} = \theta_{km} \quad \forall (k, m) \quad (\text{see Theorem 2.3}) \quad (3.12)$$

$$M_{0kmn} = 0, \quad n \geq 1. \quad (3.13)$$

Proof. By Theorem 3.1, the derivative on the left hand side of Eq.(3.1) equals

$$\frac{d}{dt} \sum_{m=0}^{j+k} e^{a_m t} \sum_{n=0}^j M_{jkmn} t^n = \sum_{m=0}^{j+k} e^{a_m t} \sum_{n=0}^j [a_m M_{jkmn} + (n+1)M_{j,k,m,n+1}] t^n.$$

Here, in order to simplify the notation, we have added some M_{jkmn} which are necessarily 0, since $n > \min(j, j+k-m)$. The right hand side of (3.1) may be written as

$$\sum_{m=0}^{j+k} e^{a_m t} \sum_{n=0}^j [a_k M_{jkmn} + b_k M_{j,k-1,m,n} + j M_{j-1,k+1,m,n}] t^n.$$

Hence,

$$M_{j,k,m,n+1} = \frac{1}{n+1} [(a_k - a_m)M_{jkmn} + R_{jkmn}]. \quad (3.14)$$

If $k = m$, then this is (3.10). If $k \neq m$, it can be verified that (3.9) is the solution of (3.14) that satisfies the condition

$$M_{j,k,m,1+j \wedge (j+k-m)} = 0$$

(which must hold by Theorem 3.1). Eq.(3.11) results from $M_{jk}(0) = 0$ for any $j \geq 1$, while (3.12)-(3.13) follow from Theorem 2.3. \square

Eqs.(3.9)-(3.12) are easy to program, and give the explicit formulas for any moment of Y_t much faster than solving the differential equations (3.1). The recursion must proceed through the values $(j, k) = (0, 0), (0, 1), (1, 0), (0, 2), (1, 1), (2, 0)$, and so on, up to the highest required moment $EY_t^j = M_{j0}(t)$. Observe that those formulas only need to be generated symbolically once. The first three moments of Y_t are:

$$\begin{aligned} EY_t &= -\frac{\bar{x}}{\alpha} - \frac{\beta}{\alpha^2} - \frac{t\beta}{\alpha} + e^{t\alpha} \left(\frac{\bar{x}}{\alpha} + \frac{\beta}{\alpha^2} \right) \\ EY_t^2 &= \frac{\bar{x}^2}{\alpha^2} + \frac{2\bar{x}\beta}{\alpha^3} + \frac{\beta^2}{\alpha^4} - \frac{\bar{x}\gamma^2}{\alpha^3} - \frac{5\beta\gamma^2}{2\alpha^4} + t \left(\frac{2\bar{x}\beta}{\alpha^2} + \frac{2\beta^2}{\alpha^3} - \frac{\beta\gamma^2}{\alpha^3} \right) + \frac{t^2\beta^2}{\alpha^2} \\ &\quad + e^{t\alpha} \left[-\frac{2\bar{x}^2}{\alpha^2} - \frac{4\bar{x}\beta}{\alpha^3} - \frac{2\beta^2}{\alpha^4} + \frac{2\beta\gamma^2}{\alpha^4} + t \left(-\frac{2\bar{x}\beta}{\alpha^2} - \frac{2\beta^2}{\alpha^3} - \frac{2\bar{x}\gamma^2}{\alpha^2} - \frac{2\beta\gamma^2}{\alpha^3} \right) \right] \\ &\quad + e^{2t\alpha} \left(\frac{\bar{x}^2}{\alpha^2} + \frac{2\bar{x}\beta}{\alpha^3} + \frac{\beta^2}{\alpha^4} + \frac{\bar{x}\gamma^2}{\alpha^3} + \frac{\beta\gamma^2}{2\alpha^4} \right) \\ EY_t^3 &= -\frac{\bar{x}^3}{\alpha^3} - \frac{3\bar{x}^2\beta}{\alpha^4} - \frac{3\bar{x}\beta^2}{\alpha^5} - \frac{\beta^3}{\alpha^6} + \frac{3\bar{x}^2\gamma^2}{\alpha^4} + \frac{21\bar{x}\beta\gamma^2}{2\alpha^5} + \frac{15\beta^2\gamma^2}{2\alpha^6} - \frac{3\bar{x}\gamma^4}{\alpha^5} - \frac{11\beta\gamma^4}{\alpha^6} \\ &\quad + t \left(-\frac{3\bar{x}^2\beta}{\alpha^3} - \frac{6\bar{x}\beta^2}{\alpha^4} - \frac{3\beta^3}{\alpha^5} + \frac{6\bar{x}\beta\gamma^2}{\alpha^4} + \frac{21\beta^2\gamma^2}{2\alpha^5} - \frac{3\beta\gamma^4}{\alpha^5} \right) \\ &\quad + t^2 \left(-\frac{3\bar{x}\beta^2}{\alpha^3} - \frac{3\beta^3}{\alpha^4} + \frac{3\beta^2\gamma^2}{\alpha^4} \right) - \frac{t^3\beta^3}{\alpha^3} \\ &\quad + e^{t\alpha} \left[\frac{3\bar{x}^3}{\alpha^3} + \frac{9\bar{x}^2\beta}{\alpha^4} + \frac{9\bar{x}\beta^2}{\alpha^5} + \frac{3\beta^3}{\alpha^6} - \frac{3\bar{x}^2\gamma^2}{\alpha^4} - \frac{33\bar{x}\beta\gamma^2}{2\alpha^5} - \frac{27\beta^2\gamma^2}{2\alpha^6} - \frac{3\bar{x}\gamma^4}{2\alpha^5} + \frac{15\beta\gamma^4}{2\alpha^6} \right] \\ &\quad + t \left(\frac{6\bar{x}^2\beta}{\alpha^3} + \frac{12\bar{x}\beta^2}{\alpha^4} + \frac{6\beta^3}{\alpha^5} + \frac{6\bar{x}^2\gamma^2}{\alpha^3} + \frac{9\bar{x}\beta\gamma^2}{\alpha^4} - \frac{3\beta^2\gamma^2}{\alpha^5} - \frac{3\bar{x}\gamma^4}{\alpha^4} - \frac{9\beta\gamma^4}{\alpha^5} \right) \end{aligned}$$

$$\begin{aligned}
 & + t^2 \left(\frac{3\bar{x}\beta^2}{\alpha^3} + \frac{3\beta^3}{\alpha^4} + \frac{6\bar{x}\beta\gamma^2}{\alpha^3} + \frac{6\beta^2\gamma^2}{\alpha^4} + \frac{3\bar{x}\gamma^4}{\alpha^3} + \frac{3\beta\gamma^4}{\alpha^4} \right) \\
 & + e^{2t\alpha} \left[-\frac{3\bar{x}^3}{\alpha^3} - \frac{9\bar{x}^2\beta}{\alpha^4} - \frac{9\bar{x}\beta^2}{\alpha^5} - \frac{3\beta^3}{\alpha^6} - \frac{3\bar{x}^2\gamma^2}{\alpha^4} + \frac{3\bar{x}\beta\gamma^2}{2\alpha^5} + \frac{9\beta^2\gamma^2}{2\alpha^6} + \frac{3\bar{x}\gamma^4}{\alpha^5} + \frac{3\beta\gamma^4}{\alpha^6} \right. \\
 & + t \left(-\frac{3\bar{x}^2\beta}{\alpha^3} - \frac{6\bar{x}\beta^2}{\alpha^4} - \frac{3\beta^3}{\alpha^5} - \frac{6\bar{x}^2\gamma^2}{\alpha^3} - \frac{15\bar{x}\beta\gamma^2}{\alpha^4} - \frac{15\beta^2\gamma^2}{2\alpha^5} - \frac{6\bar{x}\gamma^4}{\alpha^4} - \frac{3\beta\gamma^4}{\alpha^5} \right) \\
 & \left. + e^{3t\alpha} \left(\frac{\bar{x}^3}{\alpha^3} + \frac{3\bar{x}^2\beta}{\alpha^4} + \frac{3\bar{x}\beta^2}{\alpha^5} + \frac{\beta^3}{\alpha^6} + \frac{3\bar{x}^2\gamma^2}{\alpha^4} + \frac{9\bar{x}\beta\gamma^2}{2\alpha^5} + \frac{3\beta^2\gamma^2}{2\alpha^6} + \frac{3\bar{x}\gamma^4}{2\alpha^5} + \frac{\beta\gamma^4}{2\alpha^6} \right) \right].
 \end{aligned}$$

4. Laplace transform, inverse moments and density of Y_t

In this section and the next one, the symbol \mathcal{O} (“big oh”) has the following meaning: we write

$$g(s) = \mathcal{O}(h(s)) \quad \text{as } s \rightarrow \infty$$

if $|g(s)/h(s)|$ remains bounded as $s \rightarrow \infty$. In words, $g(s)$ is big oh of $h(s)$ as s tends to infinity if $|g(s)|$ is no larger than some constant times $|h(s)|$ when s is large. We write

$$g(s) = \mathcal{o}(h(s)) \quad \text{as } s \rightarrow \infty$$

(“ $g(s)$ is little oh of $h(s)$ as $s \rightarrow \infty$ ”) if

$$\lim_{s \rightarrow \infty} \frac{g(s)}{h(s)} = 0.$$

Another symbol used below is “ \sim ”, meaning “asymptotically equal to”: we write $g(s) \sim h(s)$ if

$$\lim_{s \rightarrow \infty} \frac{g(s)}{h(s)} = 1.$$

Finally, “ i ” is now the imaginary unit of the complex numbers (that is, $i^2 = -1$).

Since the Laplace transform of the integral of the squared Bessel process may be derived explicitly (see Revuz & Yor, 1999, p.445), it is not surprising that the Laplace transform of the square-root process also has an explicit expression. A quick way to find $E \exp(-sY_t)$ is to suitably modify the CIR formula for the price of a zero-coupon bond (Cox *et al.*, 1985). Suppose the short rate follows a process of type (1.1); the bond price is known to be the expectation of the exponential of minus the integral of the short rate. Multiplying the spot rate by a positive number yields another process which also satisfies (1.1), but with different parameters. Rewriting the bond price formula for this new process immediately yields the Laplace transform of Y_t . The details follow.

Suppose X is the solution of (1.1), let $s > 0$, and define

$$\tilde{X}_t = sX_t, \quad \tilde{X}_0 = \tilde{x} = s\bar{x}.$$

Then

$$\tilde{X}_t = \tilde{x} + \int_0^t (\alpha \tilde{X}_u + \tilde{\beta}) du + \int_0^t \tilde{\gamma} \sqrt{\tilde{X}_u} dW_u,$$

where $\tilde{\beta} = \beta s$ and $\tilde{\gamma} = \gamma \sqrt{s}$. The formula for the price of a zero-coupon bond maturing in t years is (Cox *et al.*, 1985)

$$\begin{aligned} \mathbb{E} e^{-sY_t} &= \mathbb{E} e^{-\int_0^t \tilde{X}_u du} \\ &= \left[\frac{2\sqrt{\alpha^2 + 2\tilde{\gamma}^2} e^{(\sqrt{\alpha^2 + 2\tilde{\gamma}^2} - \alpha)t/2}}{(\sqrt{\alpha^2 + 2\tilde{\gamma}^2} - \alpha)(e^{\sqrt{\alpha^2 + 2\tilde{\gamma}^2}t} - 1) + 2\sqrt{\alpha^2 + 2\tilde{\gamma}^2}} \right]^{\frac{2\tilde{\beta}}{\tilde{\gamma}^2}} \\ &\quad \times \exp \left[-\tilde{x} \frac{2(e^{\sqrt{\alpha^2 + 2\tilde{\gamma}^2}t} - 1)}{(\sqrt{\alpha^2 + 2\tilde{\gamma}^2} - \alpha)(e^{\sqrt{\alpha^2 + 2\tilde{\gamma}^2}t} - 1) + 2\sqrt{\alpha^2 + 2\tilde{\gamma}^2}} \right] \\ &= \left[\frac{e^{-\alpha t/2}}{\cosh(Pt/2) - \frac{\alpha}{P} \sinh(Pt/2)} \right]^{\frac{2\tilde{\beta}}{\tilde{\gamma}^2}} \exp \left[-\frac{s\tilde{x}}{P} \frac{2 \sinh(Pt/2)}{\cosh(Pt/2) - \frac{\alpha}{P} \sinh(Pt/2)} \right], \quad (4.1) \end{aligned}$$

where $P = P(s) = \sqrt{\alpha^2 + 2\tilde{\gamma}^2 s}$.

The Laplace transform of Y_t is finite in a neighborhood of the origin. This is seen by noting that

$$\cosh(Pt/2), \quad \frac{\alpha}{P} \sinh(Pt/2)$$

are both analytic functions of P , and that their difference does not vanish at $P(0) = |\alpha|$, for any values of the parameters α, β, γ considered, and for any $t > 0$. Consequently, (1) all the moments of Y_t are finite, and (2) they are given by the familiar formula

$$\mathbb{E} Y_t^k = (-1)^k \frac{d^k}{ds^k} \mathbb{E} e^{-sY_t} \Big|_{s=0}, \quad k \in \mathbb{N}.$$

The Laplace transform gives a third method of calculating the moments of Y_t . The author has compared the computation times for the moments of Y_t up to order 6 using Mathematica; the recursive method of Section 3 does a lot better than the others, especially for higher moments. As an example, on what is now a rather slow machine (Macintosh 3400 180 Mhz), Mathematica took 377 seconds to find the general symbolic formula for the 6th moment by differentiating the Laplace transform, and a bit more to find the general formulas for moments of order 1 to 6 by solving differential equations (3.1). The recursive method produced all 6 moments in a little more than 8 seconds. A program recursively calculating just the numerical values of the moments (rather than the general symbolic formulas) would of course execute much faster, especially in C.

The next theorem concerns the moments and MGF of $1/Y_t$.

Theorem 4.1. (a) $\mathbb{E} Y_t^r$ is finite for all $r \in \mathbb{R}$, and

$$\mathbb{E} Y_t^{-q} = \frac{1}{\Gamma(q)} \int_0^\infty s^{q-1} \mathbb{E} \exp(-sY_t) ds, \quad q > 0.$$

The integrated square-root process

$$(b) \mathbb{E} \exp\left(\frac{p}{Y_t}\right) = 1 + \sqrt{p} \int_0^\infty s^{-\frac{1}{2}} I_1(2\sqrt{ps}) \mathbb{E} e^{-sY_t} ds \quad \text{for all } p \geq 0.$$

This is finite if, and only if, $0 \leq p < \frac{1}{2\gamma^2} (\beta t + \bar{x})^2$.

$$(c) \mathbb{E} \exp\left(p \int_0^t \frac{1}{X_s} ds\right) = \infty \quad \text{for all } p \geq \frac{1}{2\gamma^2} \left(\beta + \frac{\bar{x}}{t}\right)^2.$$

Proof. By (2.14),

$$P = \gamma\sqrt{2s} \sqrt{1 + \frac{\alpha^2}{2\gamma^2 s}} = \gamma\sqrt{2s} [1 + \mathcal{O}(s^{-1})].$$

From this, we can find the asymptotic behaviour of the first factor in (4.1):

$$\begin{aligned} \left[\frac{e^{-\alpha t/2}}{\cosh(Pt/2) - \frac{\alpha}{P} \sinh(Pt/2)} \right]^{\frac{2\beta}{\gamma^2}} &= 2^{\frac{2\beta}{\gamma^2}} e^{-\alpha\beta t/\gamma^2} \left[\left(1 - \frac{\alpha}{P}\right) e^{Pt/2} + \left(1 + \frac{\alpha}{P}\right) e^{-Pt/2} \right]^{-\frac{2\beta}{\gamma^2}} \\ &= 2^{\frac{2\beta}{\gamma^2}} e^{-\alpha\beta t/\gamma^2 - \beta Pt/\gamma^2} \left[\left(1 - \frac{\alpha}{P}\right) + \left(1 + \frac{\alpha}{P}\right) e^{-Pt} \right]^{-\frac{2\beta}{\gamma^2}} \\ &\sim 2^{\frac{2\beta}{\gamma^2}} e^{-\alpha\beta t/\gamma^2 - \beta t\sqrt{2s}/\gamma} \end{aligned}$$

as s tends to infinity. Now turn to the second factor:

$$\begin{aligned} -\frac{s\bar{x}}{P} \frac{2 \sinh(Pt/2)}{\cosh(Pt/2) - \frac{\alpha}{P} \sinh(Pt/2)} &= -\frac{2s\bar{x}}{\sqrt{\alpha^2 + 2\gamma^2 s}} \frac{1}{\coth(Pt/2) - \frac{\alpha}{\sqrt{\alpha^2 + 2\gamma^2 s}}} \\ &= -\frac{\sqrt{2s\bar{x}}}{\gamma} \left(1 + \frac{\alpha^2}{2\gamma^2 s}\right)^{-\frac{1}{2}} \frac{1}{\frac{1+e^{-Pt}}{1-e^{-Pt}} - \frac{\alpha}{\gamma\sqrt{2s}} \left(1 + \frac{\alpha^2}{2\gamma^2 s}\right)^{-\frac{1}{2}}} \\ &= -\frac{\sqrt{2s\bar{x}}}{\gamma} [1 + \mathcal{O}(s^{-1})] \frac{1}{1 + \mathcal{O}(e^{-Pt}) - \frac{\alpha}{\gamma\sqrt{2s}} + \mathcal{O}(s^{-\frac{3}{2}})} \\ &= -\frac{\sqrt{2s\bar{x}}}{\gamma} [1 + \mathcal{O}(s^{-1})] \left[1 + \frac{\alpha}{\gamma\sqrt{2s}} + \mathcal{O}(s^{-1})\right] \\ &= -\frac{\sqrt{2s\bar{x}}}{\gamma} - \frac{\alpha\bar{x}}{\gamma^2} + \mathcal{O}(s^{-\frac{1}{2}}). \end{aligned}$$

From these expressions we get

$$\mathbb{E} e^{-sY_t} \sim 2^{\frac{2\beta}{\gamma^2}} \exp\left\{ -(\beta t + \bar{x}) \left(\frac{\alpha}{\gamma^2} + \frac{\sqrt{2s}}{\gamma} \right) \right\} \quad \text{as } s \rightarrow \infty. \quad (4.2)$$

By this asymptotic identity and (2.18), $E Y_t^{-q}$ is finite if, and only if, the integral

$$2^{\frac{2q}{\gamma^2}} e^{-\frac{q}{\gamma^2}(\beta t + \bar{x})} \frac{1}{\Gamma(q)} \int_0^\infty s^{q-1} \exp \left\{ -\frac{\sqrt{2s}}{\gamma} (\beta t + \bar{x}) \right\} ds,$$

is finite, which is true for all $q > 0$. This proves part (a) of the theorem.

Now turn to part (b). For $p \geq 0$,

$$\begin{aligned} E \exp \left(\frac{p}{Y_t} \right) &= \sum_{n=0}^{\infty} \frac{p^n}{n!} E Y_t^{-n} \\ &= 1 + \int_0^\infty \sum_{n=1}^{\infty} \frac{p^n s^{n-1}}{\Gamma(n)\Gamma(n+1)} E \exp(-s Y_t) ds \\ &= 1 + \int_0^\infty \sqrt{\frac{p}{s}} \sum_{k=0}^{\infty} \frac{(2\sqrt{ps}/2)^{1+2k}}{\Gamma(k+1)\Gamma(k+2)} E \exp(-s Y_t) ds \\ &= 1 + \int_0^\infty \sqrt{\frac{p}{s}} I_1(2\sqrt{ps}) E \exp(-s Y_t) ds \end{aligned}$$

(see the definition of the modified Bessel function I_ν in (2.16)).

It remains to be seen for what $p > 0$ the above integral is finite. The integrand is continuous in s , and bounded near the origin. It is known that

$$I_\nu(x) \sim \frac{e^x}{\sqrt{2\pi x}}, \quad x \rightarrow \infty \quad (4.3)$$

(Lebedev, 1972, p.123). Combining this with (4.2), we find that

$$\sqrt{\frac{p}{s}} I_1(2\sqrt{ps}) E \exp(-s Y_t) \sim C s^{-3/4} \exp \left\{ \sqrt{2s} \left[\sqrt{2p} - \frac{1}{\gamma} (\beta t + \bar{x}) \right] \right\}$$

as $s \rightarrow \infty$ (here $C > 0$ depends on p but not on s). The function on the right is integrable over $[1, \infty)$ if, and only if,

$$p < \frac{1}{2\gamma^2} (\beta t + \bar{x})^2.$$

To prove (c), note that by the Cauchy-Schwarz inequality

$$\left[\int_0^t d\tau \right]^2 \leq \int_0^t \frac{d\tau}{X_\tau} \int_0^t X_\tau d\tau,$$

and thus

$$p \int_0^t \frac{d\tau}{X_\tau} \geq \frac{pt^2}{\int_0^t X_\tau d\tau}.$$

The result follows from (b). □

By contrast, it is easy to see that the density of X_t behaves like $x^{\bar{\nu}-1}$ as $x \downarrow 0$, with $\bar{\nu} = 2\beta/\gamma^2$ (see (2.17) and Theorem 2.4), and thus $\mathbb{E} X_t^p < \infty$ for $p > -\bar{\nu}$ only, whatever the initial condition $\bar{x} \geq 0$, and, consequently, $\mathbb{E} \exp(q/X_t) = \infty$ for any $q > 0$. Thus, the density of Y_t tends to 0 much quicker than that of X_t , as the argument tends to 0. This is a little unexpected, especially since $Y_0 = 0$ with probability one, irrespective of $X_0 = \bar{x}$. An intuitive explanation is that Y_t is close to 0 if X_s is close to 0 for “many” s between 0 and t , which is less likely than X_s being close to 0 for one particular s ; integration is therefore a “smoothing” operation in this case. This relates to Theorem 4.2 below, which says that the density of Y_t is “infinitely flat” at the origin (that is, all its derivatives vanish).

Part (c) of the theorem is related to absence of arbitrage and changes of measures in the financial model

$$dS = \mu S dt + \sqrt{X} S dV,$$

where X satisfies (1.1) and (V, W) is two-dimensional Brownian motion (possibly correlated). If one attempts the usual Girsanov change of measure to make $\{e^{-rt} S_t\}$ a martingale, then one has the problem of checking that the stochastic exponential of minus the integral of the market price of risk

$$\frac{\mu - r}{\sqrt{X}}$$

has expectation equal to one (which would make it a martingale, see Karatzas & Shreve (1998), p.12). A sufficient condition for this to hold is Novikov’s condition

$$\mathbb{E} \exp\left(\frac{1}{2} \int_0^t \frac{(\mu - r)^2}{X_s} ds\right) < \infty.$$

Theorem 4.1 shows that Novikov’s condition fails if

$$(\mu - r)^2 \geq \frac{1}{\gamma^2} \left(\beta + \frac{\bar{x}}{t}\right)^2$$

(but does not say that the condition holds otherwise).

Observe that, by Theorem 2.4, $\mathbb{E}(1/X_s) = \infty$ for all $s > 0$ if $2\beta/\gamma^2 \leq 1$, which implies

$$\mathbb{E} \int_0^t \frac{1}{X_s} ds = \infty,$$

and thus $\mathbb{E} e^{p \int_0^t ds/X_s} = \infty$ for all $p > 0$. More generally, since $\mathbb{E} X_t^{-q} = \infty$ for all $q \geq 2\beta/\gamma^2$, it is plausible that not all moments of $\int_0^t \frac{1}{X_s} ds$ are finite (though the author has not been able to prove this), which would then imply $\mathbb{E} e^{p \int_0^t ds/X_s} = \infty$ for all $p > 0$, for any β, γ and $t > 0$.

Theorem 4.2. *Suppose $t > 0$ and let $g_t(\cdot)$ be the density of Y_t with respect to Lebesgue measure on \mathbb{R} . Then $g_t(x)$ exists and is a continuous, uniformly bounded function of $x \in \mathbb{R}$. The same applies to $(d^n/dx^n)g_t(x)$, for any $n \geq 1$.*

Proof. Apply the following well-known property of characteristic functions (see Theorem 3 and Lemma 2 in Feller (1971), pp.509 and 512): *Suppose a r.v. U satisfies*

$$\int_{-\infty}^{\infty} |\mathbb{E} e^{i\zeta U \zeta^n}| d\zeta < \infty$$

for some $n \geq 1$; then the distribution of U has a density which is everywhere continuous and uniformly bounded, and the n th derivative of this density is also everywhere continuous and uniformly bounded.

The characteristic function of Y_t is

$$\begin{aligned} \psi(\zeta) &= \mathbb{E} e^{i\zeta Y_t} = \mathbb{E} e^{-s Y_t} \Big|_{s=-i\zeta} \\ &= \left[\frac{e^{-\alpha t/2}}{\cosh(Zt/2) - \frac{\alpha}{Z} \sinh(Zt/2)} \right]^{\frac{2\beta}{\gamma^2}} \exp \left[\frac{i\zeta \bar{x}}{Z} \frac{2 \sinh(Zt/2)}{\cosh(Zt/2) - \frac{\alpha}{Z} \sinh(Zt/2)} \right], \end{aligned}$$

where $Z = Z(\zeta) = \sqrt{\alpha^2 - 2\gamma^2 \zeta i}$. First, consider $\psi(\zeta)$ as $\zeta \rightarrow \infty$. We have (see (2.14))

$$\begin{aligned} Z(\zeta) - \gamma(1-i)\sqrt{\zeta} &= \gamma(1-i)\sqrt{\zeta} \left(\sqrt{1 - \frac{\alpha^2}{2\gamma^2 \zeta i}} - 1 \right) \\ &= \gamma(1-i)\sqrt{\zeta} \sum_{k=1}^{\infty} \left(\frac{\alpha^2}{2\gamma^2 \zeta i} \right)^k \frac{(-\frac{1}{2})_k}{k!} \\ &\rightarrow 0 \end{aligned}$$

as $\zeta \rightarrow \infty$. This implies

$$\begin{aligned} e^{Zt/2} &\sim e^{\gamma(1-i)\sqrt{\zeta}t/2} \\ \cosh(Zt/2) - \frac{\alpha}{Z} \sinh(Zt/2) &\sim \frac{1}{2} e^{\gamma(1-i)\sqrt{\zeta}t/2}. \end{aligned}$$

Moreover,

$$\begin{aligned} \frac{i\zeta \bar{x}}{Z} \frac{2 \sinh(Zt/2)}{\cosh(Zt/2) - \frac{\alpha}{Z} \sinh(Zt/2)} &= \frac{2i\zeta \bar{x}}{Z} \frac{1}{\coth(Zt/2) - \frac{\alpha}{Z}} \\ &= \frac{(-1+i)\bar{x}\sqrt{\zeta}}{\gamma} \left(1 - \frac{\alpha^2}{2\gamma^2 \zeta i} \right)^{-\frac{1}{2}} \frac{1}{\frac{1+e^{-Zt}}{1-e^{-Zt}} - \frac{\alpha}{\gamma(1-i)\sqrt{\zeta}} \left(1 - \frac{\alpha^2}{2\gamma^2 \zeta i} \right)^{-\frac{1}{2}}} \end{aligned}$$

The integrated square-root process

$$\begin{aligned}
&= \frac{(-1+i)\bar{x}\sqrt{\zeta}}{\gamma} (1 + \mathcal{O}(\zeta^{-1})) \frac{1}{1 + \mathcal{O}(e^{-Zt}) - \frac{\alpha}{\gamma(1-i)\sqrt{\zeta}}(1 + \mathcal{O}(\zeta^{-1}))} \\
&= \frac{(-1+i)\bar{x}\sqrt{\zeta}}{\gamma} (1 + \mathcal{O}(\zeta^{-1})) \left(1 + \frac{\alpha}{\gamma(1-i)\sqrt{\zeta}} + \mathcal{O}(\zeta^{-1}) \right) \\
&= \frac{(-1+i)\bar{x}\sqrt{\zeta}}{\gamma} - \frac{\alpha\bar{x}}{\gamma^2} + \mathcal{O}(\zeta^{-\frac{1}{2}})
\end{aligned}$$

Putting all this together, we find

$$|\psi(\zeta)| \sim C_1 e^{-C_2 \sqrt{\zeta}}$$

as $\zeta \rightarrow \infty$, where $C_1, C_2 > 0$ do not depend on ζ . Hence

$$\int_0^\infty |\psi(\zeta)\zeta^n| d\zeta < \infty$$

for any $n \geq 0$. The integral from $-\infty$ to 0 is handled similarly: as $\zeta \rightarrow -\infty$,

$$\begin{aligned}
Z(\zeta) - \gamma(1+i)\sqrt{-\zeta} &\rightarrow 0, \\
e^{Zt/2} &\sim e^{\gamma(1+i)\sqrt{-\zeta}t/2} \\
\cosh(Zt/2) - \frac{\alpha}{Z} \sinh(Zt/2) &\sim \frac{1}{2} e^{\gamma(1+i)\sqrt{-\zeta}t/2} \\
\frac{i\zeta\bar{x}}{Z} \frac{2 \sinh(Zt/2)}{\cosh(Zt/2) - \frac{\alpha}{Z} \sinh(Zt/2)} &= -\frac{(1+i)\bar{x}\sqrt{-\zeta}}{\gamma} - \frac{\alpha\bar{x}}{\gamma^2} + \mathcal{O}((-\zeta)^{-\frac{1}{2}}) \\
|\psi(\zeta)| &\sim C_3 e^{-C_4 \sqrt{-\zeta}}. \quad \square
\end{aligned}$$

5. Relationships between the Laplace transforms of U and $1/U$

The formula in Theorem 4.1 (b) applies for any random variable $U > 0$. The same proof leads to the identity

$$\mathbb{E} e^{p/U} = 1 + \int_0^\infty \sqrt{\frac{p}{s}} I_1(2\sqrt{ps}) \mathbb{E} \exp(-sU) ds, \quad p \geq 0 \quad (5.1)$$

(both sides are simultaneously finite or infinite, by Fubini's Theorem). Formula (5.1) strangely resembles a known relationship for Laplace transforms (Oberhettinger & Badii, 1973, p.4):

$$\int_0^\infty e^{-pt} t^{\nu-1} f(t^{-1}) dt = \int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \left[\int_0^\infty e^{-st} f(t) dt \right] ds, \quad \nu > -1. \quad (5.2)$$

Here, J_ν is the ordinary Bessel function of order ν (Lebedev, 1972, p.102),

$$J_\nu(z) = \sum_{k=0}^{\infty} \frac{(-1)^k (z/2)^{\nu+2k}}{\Gamma(k+1)\Gamma(k+\nu+1)}, \quad |\arg z| < \pi, \quad \nu \in \mathbb{C}.$$

Let us rephrase (5.2) in probabilistic terms, by considering that $f(\cdot)$ is the density of a random variable $U > 0$:

$$\mathbb{E}U^{-\nu-1}e^{-p/U} = \int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \mathbb{E}e^{-sU} ds, \quad \nu > -1. \quad (5.3)$$

(Oberhettinger & Badii do not specify for which p this holds, but this is clarified in the theorem below.) Observe that $\nu = -1$ is not allowed here, so that (5.3) does not apply to $\mathbb{E}e^{-p/U}$. We now derive some more general formulas, which have (5.1) and (5.3) as particular cases.

Theorem 5.1. *Suppose $p > 0$, $r \in \mathbb{R}$ and $U > 0$ a.s.*

(a) *If $\nu > -1$, then*

$$\mathbb{E}U^{r-\nu-1}e^{p/U} = \int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} I_\nu(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds.$$

Both sides may be infinite.

(b) *If $\nu \leq -1$, then*

$$\begin{aligned} \mathbb{E}U^{r-\nu-1}e^{p/U} &= \sum_{0 \leq n \leq -\nu-1} \frac{p^n}{n!} \mathbb{E}U^{r-\nu-n-1} \\ &+ \int_0^\infty \left[\left(\frac{s}{p}\right)^{\frac{\nu}{2}} I_\nu(2\sqrt{ps}) - \sum_{0 \leq n \leq -\nu-1} \frac{p^n s^{n+\nu}}{n! \Gamma(n+\nu+1)} \right] \mathbb{E}U^r e^{-sU} ds. \end{aligned}$$

Both sides may be infinite. If, moreover, $-\nu-1 \in \mathbb{N}$, then

$$\mathbb{E}U^{r-\nu-1}e^{p/U} = \sum_{0 \leq n \leq -\nu-1} \frac{p^n}{n!} \mathbb{E}U^{r-\nu-n-1} + \int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} I_{-\nu}(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds.$$

(c) *If $\nu > -1$, $\mathbb{E}U^r < \infty$, $\mathbb{E}U^{r-\nu-1} < \infty$, then*

$$\mathbb{E}U^{r-\nu-1}e^{-p/U} = \int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds.$$

Both sides are finite, and the integral on the right may be improper (that is, it may not converge absolutely) if $\nu < -1/2$.

(d) If $\nu \leq -1$, $EU^r < \infty$, $EU^{r-\nu-1} < \infty$, then

$$EU^{r-\nu-1}e^{-p/U} = \sum_{0 \leq n \leq -\nu-1} \frac{(-p)^n}{n!} EU^{r-\nu-n-1} + \int_0^\infty \left[\left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) - \sum_{0 \leq n \leq -\nu-1} \frac{(-p)^n s^{n+\nu}}{n! \Gamma(n+\nu+1)} \right] EU^r e^{-sU} ds.$$

Both sides are finite, and the integral on the right may be improper. If, moreover, $-\nu-1 \in \mathbb{N}$, then

$$EU^{r-\nu-1}e^{-p/U} = \sum_{0 \leq n \leq -\nu-1} \frac{(-p)^n}{n!} EU^{r-\nu-n-1} + (-1)^\nu \int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_{-\nu}(2\sqrt{ps}) EU^r e^{-sU} ds.$$

Proof. (a) Because all the terms in the following series are positive, the equalities hold whether the resulting expressions are finite or infinite:

$$\begin{aligned} EU^{r-\nu-1}e^{p/U} &= \sum_{n=0}^{\infty} \frac{p^n}{n!} EU^{r-\nu-n-1} \\ &= \sum_{n=0}^{\infty} \frac{p^n}{n!} EU^r \int_0^\infty \frac{s^{n+\nu}}{\Gamma(\nu+n+1)} e^{-sU} ds \\ &= \int_0^\infty \left(\sum_{n=0}^{\infty} \frac{p^n}{n!} \frac{s^{n+\nu}}{\Gamma(\nu+n+1)} \right) EU^r e^{-sU} ds \\ &= \int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} I_\nu(2\sqrt{ps}) EU^r e^{-sU} ds. \end{aligned}$$

(b) The first formula results from

$$EU^{r-\nu-1}e^{p/U} = \sum_{0 \leq n \leq -\nu-1} \frac{p^n}{n!} EU^{r-\nu-n-1} + \int_0^\infty \sum_{n > -\nu-1} \frac{p^n s^{n+\nu}}{n! \Gamma(\nu+n+1)} EU^r e^{-sU} ds.$$

The second one then follows from (Lebedev, 1972, p.110)

$$\frac{1}{\Gamma(-m)} = 0, \quad I_{-m}(z) = I_m(z), \quad m \in \mathbb{N}.$$

(c) Let $U_\epsilon = \max(U, \epsilon)$ for $\epsilon \geq 0$ (whence $U_0 = U$). For $\epsilon > 0$, $1/U_\epsilon$ is bounded below by 0, and above by $1/\epsilon$, whence

$$EU^r U_\epsilon^{-\nu-1} e^{p/U_\epsilon} < \infty.$$

Therefore the series below converge absolutely, and so it is permitted to reverse the order of summation, integration and expectation at will:

$$\begin{aligned} \mathbb{E}U^r U_\epsilon^{-\nu-1} e^{-p/U_\epsilon} &= \int_0^\infty \left(\sum_{n=0}^\infty \frac{(-p)^n}{n!} \frac{s^{n+\nu}}{\Gamma(\nu+n+1)} \right) \mathbb{E}U^r e^{-sU_\epsilon} ds \\ &= \int_0^\infty \left(\frac{s}{p} \right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \mathbb{E}U^r e^{-sU_\epsilon} ds. \end{aligned} \quad (5.4)$$

Now $-\nu - 1 < 0$, so

$$0 \leq U^r U_\epsilon^{-\nu-1} e^{-p/U_\epsilon} \leq CU^r$$

where $C = \sup_{x>0} x^{\nu+1} e^{-px} < \infty$, and U^r is integrable by assumption. Thus

$$\mathbb{E}U^r U_\epsilon^{-\nu-1} e^{-p/U_\epsilon} \rightarrow \mathbb{E}U^{r-\nu-1} e^{-p/U}$$

as $\epsilon \rightarrow 0+$. Next, turn to the right hand side of (5.4).

Case 1: $\nu \geq -\frac{1}{2}$

The asymptotic expression (Lebedev, 1972, p.122)

$$J_\nu(x) = \left(\frac{2}{\pi x} \right)^{\frac{1}{2}} \cos\left(x - (2\nu + 1)\frac{\pi}{4}\right) + \mathcal{O}(x^{-3/2}) \quad \text{as } x \rightarrow \infty$$

implies that there is $C > 0$ such that

$$|J_\nu(x)| \leq Cx^{-\frac{1}{2}}, \quad x > 0.$$

Therefore

$$\left| \left(\frac{s}{p} \right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \mathbb{E}U^r e^{-sU_\epsilon} \right| \leq C_1 s^{\frac{\nu}{2}-\frac{1}{4}} \mathbb{E}U^r e^{-sU}.$$

Now $\nu \geq -\frac{1}{2}$ implies $\frac{\nu}{2} + \frac{3}{4} > 0$, and so

$$\int_0^\infty s^{\frac{\nu}{2}-\frac{1}{4}} \mathbb{E}U^r e^{-sU} ds = \mathbb{E}U^{r-\frac{\nu}{2}-\frac{3}{4}} \Gamma\left(\frac{\nu}{2} + \frac{3}{4}\right).$$

This is finite, because $r - \nu - 1 \leq r - \frac{\nu}{2} - \frac{3}{4} \leq r$, and it is assumed that both $\mathbb{E}U^r$ and $\mathbb{E}U^{r-\nu-1}$ are finite. Finally, the right hand side of (5.4) converges to

$$\int_0^\infty \left(\frac{s}{p} \right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds$$

as $\epsilon \rightarrow 0+$, by dominated convergence.

Case 2: $\nu \in (-1, -\frac{1}{2})$ The above arguments break down if $\nu < -\frac{1}{2}$, because then

$$r - \frac{\nu}{2} - \frac{3}{4} < r - \nu - 1.$$

The problem can be solved either by assuming that $\mathbb{E}U^{r-\frac{\nu}{2}-\frac{3}{4}} < \infty$ or else as follows.

Write the integral on the right hand side of (5.4) as $\int_0^1 + \int_1^\infty$. It is immediately seen that

$$\int_0^1 \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \mathbb{E}U^r e^{-sU_\epsilon} ds \rightarrow \int_0^1 \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds$$

as $\epsilon \rightarrow 0+$, by dominated convergence. For the second integral, consider separately each term on the right of

$$J_\nu(2\sqrt{ps}) = C_2 s^{-\frac{1}{4}} \cos(2\sqrt{ps} - (2\nu + 1)\frac{\pi}{4}) + R(s) \quad \text{as } s \rightarrow \infty$$

where $C_2 > 0$ and $R(s) = \mathcal{O}(s^{-\frac{3}{4}})$. Since $\frac{\nu}{2} - \frac{3}{4} < -1$, the integral

$$\int_1^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} R(s) \mathbb{E}U^r e^{-sU_\epsilon} ds$$

converges absolutely for $\epsilon \geq 0$, and so convergence as $\epsilon \rightarrow 0+$ to

$$\int_1^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} R(s) \mathbb{E}U^r e^{-sU} ds$$

follows at once. The only remaining problem is to show that

$$\int_0^\infty s^{\frac{\nu}{2}-\frac{1}{4}} \cos(2\sqrt{ps} - (2\nu + 1)\frac{\pi}{4}) \mathbb{E}U^r e^{-sU_\epsilon} ds$$

converges to the same expression with $\epsilon = 0$, as $\epsilon \rightarrow 0+$. Perform the change of variable $u = 2\sqrt{ps}$, $s = u^2/(4p)$, to obtain

$$\begin{aligned} & 2 \int_{2\sqrt{p}}^\infty \left(\frac{u^2}{4p}\right)^{\frac{\nu}{2}-\frac{1}{4}} u \cos(u - (2\nu + 1)\frac{\pi}{4}) \mathbb{E}U^r e^{-u^2 U_\epsilon/(4p)} du \\ &= C_3 \int_{2\sqrt{p}}^\infty u^{\nu+\frac{1}{2}} \cos(u - (2\nu + 1)\frac{\pi}{4}) \mathbb{E}U^r e^{-u^2 U_\epsilon/(4p)} du \\ &= C_3 \left\{ u^{\nu+\frac{1}{2}} \sin(u - (2\nu + 1)\frac{\pi}{4}) \mathbb{E}U^r e^{-u^2 U_\epsilon/(4p)} \Big|_{u=2\sqrt{p}}^{u \rightarrow \infty} \right\} \\ & \quad - C_3 \int_{2\sqrt{p}}^\infty \sin(u - (2\nu + 1)\frac{\pi}{4}) d \left[u^{\nu+\frac{1}{2}} \mathbb{E}U^r e^{-u^2 U_\epsilon/(4p)} \right]. \end{aligned} \quad (5.5)$$

Now $u^{\nu+\frac{1}{2}}$ tends to 0 as u tends to ∞ , so the expression inside the curly brackets reduces to

$$-u^{\nu+\frac{1}{2}} \sin(u - (2\nu + 1)\frac{\pi}{4}) \mathbb{E}U^r e^{-u^2 U_\epsilon / (4p)} \Big|_{u=2\sqrt{p}},$$

which poses no problem. The integral in (5.5) converges absolutely, since

$$\lim_{u \rightarrow \infty} u^{\nu+\frac{1}{2}} \mathbb{E}U^r e^{-u^2 U_\epsilon / (4p)} = 0.$$

To prove that it converges to

$$\int_{2\sqrt{p}}^{\infty} \sin(u - (2\nu + 1)\frac{\pi}{4}) d \left[u^{\nu+\frac{1}{2}} \mathbb{E}U^r e^{-u^2 U_\epsilon / (4p)} \right],$$

apply the Helly-Bray Theorem (Loève, 1977, p.184; Kolmogorov & Fomine, 1975, p.370). The functions

$$F_\epsilon(u) = u^{\nu+\frac{1}{2}} \mathbb{E}U^r e^{-u^2 U_\epsilon / (4p)}, \quad \epsilon \geq 0$$

are of bounded variation on \mathbb{R}_+ , and $\{F_\epsilon; \epsilon > 0\}$ converge completely (Loève, 1977, p.180) to F_0 as $\epsilon \rightarrow 0+$. Since $g(u) = \sin(u - (2\nu + 1)\frac{\pi}{4})$ is bounded, we conclude that

$$\int_{2\sqrt{p}}^{\infty} g dF_\epsilon \rightarrow \int_{2\sqrt{p}}^{\infty} g dF_0 \quad \text{as } \epsilon \rightarrow 0+.$$

(d) Proceed as in (b) and (c) to get

$$\begin{aligned} & \mathbb{E}U^r U_\epsilon^{-\nu-1} e^{p/U_\epsilon} \\ &= \sum_{0 \leq n \leq -\nu-1} \frac{(-p)^n}{n!} \mathbb{E}U^r U_\epsilon^{-\nu-n-1} \\ &+ \int_0^\infty \left[\left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) - \sum_{0 \leq n \leq -\nu-1} \frac{(-p)^n s^{n+\nu}}{n! \Gamma(\nu+n+1)} \right] \mathbb{E}U^r e^{-sU_\epsilon} ds \end{aligned}$$

for any $\epsilon > 0$. The first sum poses no problem. Split the integral into $\int_0^1 + \int_1^\infty$. The former tends to the right limit by dominated convergence. To prove convergence of \int_1^∞ , consider each term of the integrand separately. The first one converges to

$$\int_1^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds$$

by the arguments given in Case 2 of (c). Each of the other terms gives rise to an absolutely convergent integral, since $n + \nu < -1$, and dominated convergence does the rest.

Finally, in the special case $-\nu - 1 \in \mathbb{N}$, the sum inside the integral vanishes (as in (b)), and one applies (Lebedev, 1972, p.103)

$$J_{-n}(z) = (-1)^n J_n(z), \quad n = 1, 2, \dots \quad \square$$

Theorem 4.1(b) results from letting $\nu = -1$ in Theorem 5.1(b). This theorem has the following consequences, which relate the existence of $\mathbb{E}U^{-\nu-1}e^{p/U}$, which is itself determined by the behaviour of the distribution of U near the origin, to the behaviour of $\mathbb{E}e^{-sU}$ near infinity.

Corollary 5.2. *Suppose $p > 0$, $\nu \in \mathbb{R}$ and $U > 0$ a.s.*

(a) *If $\mathbb{E}U^{-\nu-1}e^{p/U} < \infty$, then*

$$\liminf_{s \rightarrow \infty} s^{\frac{\nu}{2} - \frac{1}{4}} e^{2\sqrt{ps}} \mathbb{E}e^{-sU} = 0.$$

(b) *If $\nu > -1$ and if, for some $\epsilon > 0$, $\mathbb{E}e^{-sU} = \mathcal{O}(s^{-\frac{3}{4} - \frac{\nu}{2} - \epsilon} e^{-2\sqrt{ps}})$ as $s \rightarrow \infty$, then $\mathbb{E}U^{-\nu-1}e^{p/U} < \infty$. The same holds for $\nu \leq -1$, with the additional assumption $\mathbb{E}U^{-\nu-1} < \infty$.*

Proof. (a) First suppose $\nu > -1$, and let $r = 0$ in Theorem 5.1(a). If the integral converges, then necessarily the integrand must have a limit inferior equal to 0. By (4.3),

$$\liminf_{s \rightarrow \infty} s^{\frac{\nu}{2}} I_{\nu}(2\sqrt{ps}) \mathbb{E}e^{-sU} = C \liminf_{s \rightarrow \infty} s^{\frac{\nu}{2} - \frac{1}{4}} e^{2\sqrt{ps}} \mathbb{E}e^{-sU},$$

where $C > 0$ is a constant. For $\nu \leq -1$, let $r = 0$ in Theorem 5.1(b). If the left hand side of the first formula is finite, then the integral on the other side must be finite. The situation is seen to be the same as when $\nu > -1$, once it is observed that

$$\int_1^{\infty} s^{n+\nu} \mathbb{E}e^{-sU} ds < \infty$$

for any $n < \nu + 1$.

(b) Suppose $\nu > -1$. The assumptions imply that there is a constant C such that

$$\begin{aligned} \mathbb{E}U^{-\nu-1}e^{p/U} &= \left[\int_0^1 + \int_1^{\infty} \right] \left(\frac{s}{p} \right)^{\frac{\nu}{2}} I_{\nu}(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds \\ &\leq \int_0^1 \left(\frac{s}{p} \right)^{\frac{\nu}{2}} I_{\nu}(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds + C \int_1^{\infty} s^{\frac{\nu}{2}} I_{\nu}(2\sqrt{ps}) s^{-\frac{3}{4} - \frac{\nu}{2} - \epsilon} e^{-2\sqrt{ps}} ds \\ &= \int_0^1 \left(\frac{s}{p} \right)^{\frac{\nu}{2}} I_{\nu}(2\sqrt{ps}) \mathbb{E}U^r e^{-sU} ds + C \int_1^{\infty} s^{\frac{1}{4}} e^{-2\sqrt{ps}} I_{\nu}(2\sqrt{ps}) s^{-1-\epsilon} ds \end{aligned}$$

The result follows from (4.3). If $\nu \leq -1$ and, moreover, $\mathbb{E}U^{-\nu-1} < \infty$, then

$$\mathbb{E}U^{-\nu-n-1} < \infty \quad \forall 0 \leq n \leq -\nu - 1,$$

$$\int_1^{\infty} s^{n+\nu} \mathbb{E}e^{-sU} ds < \infty \quad \forall 0 \leq n < -\nu - 1,$$

and

$$\int_1^\infty s^{\frac{\nu}{2}} I_\nu(2\sqrt{ps}) \mathbb{E} e^{-sU} ds < \infty.$$

This implies $\mathbb{E} U^{-\nu-1} e^{p/U} < \infty$, by Theorem 5.1(b). \square

Example. Macdonald's function is defined in terms of the modified Bessel function as follows (Lebedev, 1972):

$$K_\nu(z) = \frac{\pi}{2} \frac{I_{-\nu}(z) - I_\nu(z)}{\sin(\nu\pi)}, \quad |\arg z| < \pi, \quad \nu \notin \mathbb{Z},$$

$$K_n(z) = \lim_{\nu \rightarrow n} K_\nu(z), \quad n \in \mathbb{Z}.$$

It also has the integral expression

$$K_\nu(z) = \frac{1}{2} \left(\frac{z}{2}\right)^\nu \int_0^\infty e^{-t-(z^2/4t)} t^{-\nu-1} dt, \quad |\arg(z)| < \frac{\pi}{4}.$$

Clearly $K_{-\nu}(z) = K_\nu(z)$. The generalized inverse Gaussian distribution with parameters $\alpha > 0, \beta > 0, \gamma \in \mathbb{R}$ has density

$$f(x) = \left(\frac{\alpha}{\beta}\right)^{\frac{\gamma}{2}} \frac{1}{2K_\gamma(\sqrt{\alpha\beta})} x^{\gamma-1} \exp\left\{-\frac{1}{2}\left(\alpha x + \frac{\beta}{x}\right)\right\} \mathbf{1}_{(0,\infty)}(x).$$

The notation for this distribution is $\mathbf{GIG}(\alpha, \beta, \gamma)$. It is immediate that $U \sim \mathbf{GIG}(\alpha, \beta, \gamma)$ implies $(1/U) \sim \mathbf{GIG}(\beta, \alpha, -\gamma)$. Moreover, a simple calculation shows that

$$\mathbb{E} U^r e^{-sU} = \frac{\alpha^{\frac{\gamma}{2}} \beta^{\frac{r}{2}}}{(\alpha + 2s)^{\frac{\gamma+r}{2}}} \frac{K_{\gamma+r}(\sqrt{(\alpha + 2s)\beta})}{K_\gamma(\sqrt{\alpha\beta})}, \quad \operatorname{Re}(s) > -\frac{\alpha}{2}.$$

Theorem 5.1(c) then says that

$$\int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \frac{\alpha^{\frac{\gamma}{2}} \beta^{\frac{r}{2}}}{(\alpha + 2s)^{\frac{\gamma+r}{2}}} \frac{K_{\gamma+r}(\sqrt{(\alpha + 2s)\beta})}{K_\gamma(\sqrt{\alpha\beta})} ds$$

$$= \frac{\beta^{-\frac{\gamma}{2}} \alpha^{\frac{1+\nu-r}{2}}}{(\beta + 2p)^{\frac{-\gamma+1+\nu-r}{2}}} \frac{K_{1+\nu-r-\gamma}(\sqrt{(\beta + 2p)\alpha})}{K_\gamma(\sqrt{\alpha\beta})}$$

or, equivalently,

$$\int_0^\infty \left(\frac{s}{p}\right)^{\frac{\nu}{2}} J_\nu(2\sqrt{ps}) \frac{K_{\gamma+r}(\sqrt{(\alpha + 2s)\beta})}{(\alpha + 2s)^{\frac{\gamma+r}{2}}} ds$$

$$= \frac{\beta^{-\frac{\gamma+r}{2}} \alpha^{\frac{1+\nu-r-\gamma}{2}}}{(\beta + 2p)^{\frac{-\gamma+1+\nu-r}{2}}} K_{1+\nu-r-\gamma}(\sqrt{(\beta + 2p)\alpha}). \quad (5.6)$$

By performing the change of variable $x = 2\sqrt{ps}/b$, and setting

$$\alpha = \frac{b^2 y^2}{2p}, \quad \beta = \frac{2a^2 p}{b^2}, \quad \mu = \gamma + r,$$

it can be seen that (5.6) is equivalent to the formula (Lebedev, 1972, p. 134)

$$\int_0^\infty \frac{K_\mu(a\sqrt{x^2 + y^2})}{(x^2 + y^2)^{\mu/2}} J_\nu(bx) x^{\nu+1} dx = \frac{b^\nu}{a^\mu} \left(\frac{\sqrt{a^2 + b^2}}{y} \right)^{\mu-\nu-1} K_{\mu-\nu-1}(y\sqrt{a^2 + b^2}). \quad (5.7)$$

An extension of this result is obtained by applying part (d) of Theorem 5.1 to the same distribution $\mathbf{GIG}(\alpha, \beta, \gamma)$. With the same change of variable and substitutions, we get

$$\begin{aligned} & \int_0^\infty \frac{K_\mu(a\sqrt{x^2 + y^2})}{(x^2 + y^2)^{\mu/2}} \left[J_\nu(bx) - \sum_{0 \leq n < -\nu-1} \frac{(-1)^n (bx/2)^{\nu+2n}}{n! \Gamma(n + \nu + 1)} \right] x^{\nu+1} dx \\ &= \frac{b^\nu}{a^\mu} \left(\frac{\sqrt{a^2 + b^2}}{y} \right)^{\mu-\nu-1} K_{\mu-\nu-1}(y\sqrt{a^2 + b^2}) - \frac{b^\nu}{a^{\nu+1} y^{\mu-\nu-1}} \sum_{0 \leq n \leq -\nu-1} \frac{1}{n!} \left(-\frac{b^2 y}{2a} \right)^n K_{\mu-\nu-n-1}(ay). \end{aligned} \quad (5.8)$$

The Appendix shows another way of deriving the same formula.

6. Conclusion

This paper has shown how the moments of the integral of the square-root process may be computed. Of the three methods shown, the recursive method (Theorem 3.2) is the fastest to execute, and its programming poses no problem. The paper also showed some regularly properties of the density of the integral.

Further work will consider expressing the density of Y_t as an infinite Laguerre series, as was done for the density of the integral of geometric Brownian motion in Dufresne (2001). The coefficients of the Laguerre polynomials are combinations of moments of the variable, and this is where the recursive equations of Section 3 will be essential, as they generate the moments much faster than differentiating the Laplace transform. A finite number of terms of the series would then serve as an approximation for the density. The same approach may be used for options on integrated squared volatility, or for options on average interest rates.

Acknowledgments

Financial support from the Actuarial Education and Research Fund (Society of Actuaries) is gratefully acknowledged.

References

- Carr, P. (2001). Replicating variance swaps. Presented at the Risk Course on Correlation, New York, May 22, 2001.
- Courtadon, G. (1982). The pricing of options on default-free bonds. *Journal of Financial and Quantitative Analysis* **17**: 75-100.
- Cox, J.C., Ingersoll, J.E., and Ross, S.A. (1985). A theory of the term structure of interest rates. *Econometrica* **53**: 385-407.
- Dufresne, D. (1989). Weak convergence of random growth processes with applications to insurance. *Insurance: Mathematics and Economics* **8**: 187-201.
- Dufresne (1990). The distribution of a perpetuity, with applications to risk theory and pension funding. *Scand. Actuarial. J.* **1990**: 39-79.
- Dufresne (2000). Laguerre series for Asian and other options. *Mathematical Finance* **10**: 407-428.
- Feller, W. (1971). *An Introduction to Probability Theory and Its Applications, Volume 2* (2th Edition). Wiley, Chichester.
- Heston, S.L. (1993). A closed-form solution for options with stochastic volatility with application to bond and currency options. *Rev. Fin. Studies* **6**: 327-343.
- Karatzas, I., and Shreve, S.E., 1998). *Methods of Mathematical Finance*. Springer-Verlag, New York.
- Kolmogorov, A.N., and Fomine, S.V. (1975). *Introductory Real Analysis*. Dover, New York.
- Loève, M. (1977). *Probability I* (4th Edition). Springer-Verlag, New York.
- Lebedev, N.N. (1972). *Special Functions and their Applications*. Dover, New York.
- Oberhettinger, F., and Badii, L. (1973). *Tables of Laplace Transforms*. Springer-Verlag, Berlin.
- Revuz, D., and Yor, M. (1999). *Continuous Martingales and Brownian Motion (Third Edition)*. Springer-Verlag, New York.

APPENDIX

We give another proof of formula (5.8). First, we derive a slight extension of Weber's integral (Lebedev, 1972, p.132)

$$\int_0^\infty e^{-a^2x^2} J_\nu(bx)x^{\nu+1} dx = \frac{b^\nu}{(2a^2)^{\nu+1}} e^{-b^2/4a^2}, \quad a, b > 0, \quad \text{Re}(\nu) > -1.$$

Leave $a, b > 0$, but now suppose $\nu \leq -1$ (the case of complex ν with $\text{Re}(\nu) \leq -1$ remains to be explored). Consider

$$\begin{aligned} \int_0^\infty e^{-a^2x^2} \left[J_\nu(bx) - \sum_{0 \leq n \leq -\nu-1} \frac{(-1)^n (bx/2)^{\nu+2n}}{n! \Gamma(\nu+n+1)} \right] x^{\nu+1} dx \\ &= \sum_{n > -\nu-1} \frac{(-1)^n (b/2)^{\nu+2n}}{n! \Gamma(\nu+n+1)} \int_0^\infty e^{-a^2x^2} x^{2\nu+2n+1} dx \\ &= \sum_{n > -\nu-1} \frac{(-1)^n (b/2)^{\nu+2n}}{n! \Gamma(\nu+n+1)} \frac{1}{2} a^{-2\nu-2n-2} \int_0^\infty e^{-u} u^{\nu+n} du \\ &= \sum_{n > -\nu-1} \frac{(-1)^n (b/2)^{\nu+2n}}{n!} \frac{1}{2} a^{-2\nu-2n-2} \\ &= \frac{b^\nu}{(2a^2)^{\nu+1}} \left[e^{-b^2/4a^2} - \sum_{0 \leq n \leq -\nu-1} \frac{1}{n!} \left(-\frac{b^2}{4a^2} \right)^n \right]. \end{aligned}$$

From this formula, we obtain

$$\begin{aligned} \int_0^\infty \frac{K_\mu(a\sqrt{x^2+y^2})}{(x^2+y^2)^{\mu/2}} \left[J_\nu(bx) - \sum_{0 \leq n \leq -\nu-1} \frac{(-1)^n (bx/2)^{\nu+2n}}{n! \Gamma(n+\nu+1)} \right] x^{\nu+1} dx \\ &= \int_0^\infty dx \left[J_\nu(bx) - \sum_{0 \leq n \leq -\nu-1} \frac{(-1)^n (bx/2)^{\nu+2n}}{n! \Gamma(n+\nu+1)} \right] x^{\nu+1} \frac{a^\mu}{2^{\mu+1}} \int_0^\infty \frac{dt}{t^{\mu+1}} e^{-t-a^2(x^2+y^2)/4t} \\ &= \frac{a^\mu}{2^{\mu+1}} \int_0^\infty \frac{dt}{t^{\mu+1}} e^{-t-a^2y^2/4t} \int_0^\infty dx e^{-a^2x^2/4t} \left[J_\nu(bx) - \sum_{0 \leq n \leq -\nu-1} \frac{(-1)^n (bx/2)^{\nu+2n}}{n! \Gamma(n+\nu+1)} \right] x^{\nu+1} \\ &= \frac{a^\mu}{2^{\mu+1}} \int_0^\infty \frac{dt}{t^{\mu+1}} e^{-t-a^2y^2/4t} \frac{b^\nu}{(a^2/2t)^{\nu+1}} \left[e^{-b^2t/a^2} - \sum_{0 \leq n \leq -\nu-1} \frac{1}{n!} \left(-\frac{b^2t}{a^2} \right)^n \right] \\ &= 2^{\nu-\mu} a^{\mu-2\nu-2} b^\nu \left[\int_0^\infty \frac{dt}{t^{\mu-\nu}} e^{-t-a^2y^2/4t-b^2t/a^2} - \sum_{0 \leq n \leq -\nu-1} \frac{1}{n!} \left(-\frac{b^2}{a^2} \right)^n \int_0^\infty \frac{dt}{t^{\mu-\nu-n}} e^{-t-a^2y^2/4t} \right] \\ &= \frac{b^\nu}{a^\mu} \left(\frac{\sqrt{a^2+b^2}}{y} \right)^{\mu-\nu-1} K_{\mu-\nu-1}(y\sqrt{a^2+b^2}) - \frac{b^\nu}{a^{\nu+1}y^{\mu-\nu-1}} \sum_{0 \leq n \leq -\nu-1} \frac{1}{n!} \left(-\frac{b^2y}{2a} \right)^n K_{\mu-\nu-n-1}(ay). \end{aligned}$$

The integrated square-root process

Daniel Dufresne
Department of Mathematics and Statistics
University of Montreal P.O. Box 6128, Downtown Station
Montreal
Canada H3C 3J7
Tel.: (514) 343 2315
FAX:(514) 343 5700
dufresne@dms.umontreal.ca

RESEARCH PAPER SERIES

No.	Date	Subject	Author
1	MAR 1993	AUSTRALIAN SUPERANNUATION: THE FACTS, THE FICTION, THE FUTURE	David M Knox
2	APR 1993	AN EXPONENTIAL BOUND FOR RUIN PROBABILITIES	David C M Dickson
3	APR 1993	SOME COMMENTS ON THE COMPOUND BINOMIAL MODEL	David C M Dickson
4	AUG 1993	RUIN PROBLEMS AND DUAL EVENTS	David C M Dickson Alfredo D Egídio dos Reis
5	SEP 1993	CONTEMPORARY ISSUES IN AUSTRALIAN SUPERANNUATION – A CONFERENCE SUMMARY	David M Knox John Piggott
6	SEP 1993	AN ANALYSIS OF THE EQUITY INVESTMENTS OF AUSTRALIAN SUPERANNUATION FUNDS	David M Knox
7	OCT 1993	A CRITIQUE OF DEFINED CONTRIBUTION USING A SIMULATION APPROACH	David M Knox
8	JAN 1994	REINSURANCE AND RUIN	David C M Dickson Howard R Waters
9	MAR 1994	LIFETIME INSURANCE, TAXATION, EXPENDITURE AND SUPERANNUATION (LITES): A LIFE-CYCLE SIMULATION MODEL	Margaret E Atkinson John Creedy David M Knox
10	FEB 1994	SUPERANNUATION FUNDS AND THE PROVISION OF DEVELOPMENT/VENTURE CAPITAL: THE PERFECT MATCH? YES OR NO	David M Knox
11	JUNE 1994	RUIN PROBLEMS: SIMULATION OR CALCULATION?	David C M Dickson Howard R Waters
12	JUNE 1994	THE RELATIONSHIP BETWEEN THE AGE PENSION AND SUPERANNUATION BENEFITS, PARTICULARLY FOR WOMEN	David M Knox
13	JUNE 1994	THE COST AND EQUITY IMPLICATIONS OF THE INSTITUTE OF ACTUARIES OF AUSTRALIA PROPOSED RETIREMENT INCOMES STRATEGY	Margaret E Atkinson John Creedy David M Knox Chris Haberecht
14	SEPT 1994	PROBLEMS AND PROSPECTS FOR THE LIFE INSURANCE AND PENSIONS SECTOR IN INDONESIA	Catherine Prime David M Knox

No.	Date	Subject	Author
15	OCT 1994	PRESENT PROBLEMS AND PROSPECTIVE PRESSURES IN AUSTRALIA'S SUPERANNUATION SYSTEM	David M Knox
16	DEC 1994	PLANNING RETIREMENT INCOME IN AUSTRALIA: ROUTES THROUGH THE MAZE	Margaret E Atkinson John Creedy David M Knox
17	JAN 1995	ON THE DISTRIBUTION OF THE DURATION OF NEGATIVE SURPLUS	David C M Dickson Alfredo D Egídio dos Reis
18	FEB 1995	OUTSTANDING CLAIM LIABILITIES: ARE THEY PREDICTABLE?	Ben Zehnwrith
19	MAY 1995	SOME STABLE ALGORITHMS IN RUIN THEORY AND THEIR APPLICATIONS	David C M Dickson Alfredo D Egídio dos Reis Howard R Waters
20	JUNE 1995	SOME FINANCIAL CONSEQUENCES OF THE SIZE OF AUSTRALIA'S SUPERANNUATION INDUSTRY IN THE NEXT THREE DECADES	David M Knox
21	JUNE 1995	MODELLING OPTIMAL RETIREMENT IN DECISIONS IN AUSTRALIA	Margaret E Atkinson John Creedy
22	JUNE 1995	AN EQUITY ANALYSIS OF SOME RADICAL SUGGESTIONS FOR AUSTRALIA'S RETIREMENT INCOME SYSTEM	Margaret E Atkinson John Creedy David M Knox
23	SEP 1995	EARLY RETIREMENT AND THE OPTIMAL RETIREMENT AGE	Angela Ryan
24	OCT 1995	APPROXIMATE CALCULATIONS OF MOMENTS OF RUIN RELATED DISTRIBUTIONS	David C M Dickson
25	DEC 1995	CONTEMPORARY ISSUES IN THE ONGOING REFORM OF THE AUSTRALIAN RETIREMENT INCOME SYSTEM	David M Knox
26	FEB 1996	THE CHOICE OF EARLY RETIREMENT AGE AND THE AUSTRALIAN SUPERANNUATION SYSTEM	Margaret E Atkinson John Creedy
27	FEB 1996	PREDICTIVE AGGREGATE CLAIMS DISTRIBUTIONS	David C M Dickson Ben Zehnwrith
28	FEB 1996	THE AUSTRALIAN GOVERNMENT SUPERANNUATION CO-CONTRIBUTIONS: ANALYSIS AND COMPARISON	Margaret E Atkinson
29	MAR 1996	A SURVEY OF VALUATION ASSUMPTIONS AND FUNDING METHODS USED BY AUSTRALIAN ACTUARIES IN DEFINED BENEFIT SUPERANNUATION FUND VALUATIONS	Des Welch Shauna Ferris

No.	Date	Subject	Author
30	MAR 1996	THE EFFECT OF INTEREST ON NEGATIVE SURPLUS	David C M Dickson Alfredo D Egídio dos Reis
31	MAR 1996	RESERVING CONSECUTIVE LAYERS OF INWARDS EXCESS-OFF-LOSS REINSURANCE	Greg Taylor
32	AUG 1996	EFFECTIVE AND ETHICAL INSTITUTIONAL INVESTMENT	Anthony Asher
33	AUG 1996	STOCHASTIC INVESTMENT MODELS: UNIT ROOTS, COINTEGRATION, STATE SPACE AND GARCH MODELS FOR AUSTRALIA	Michael Sherris Leanna Tedesco Ben Zehnwrith
34	AUG 1996	THREE POWERFUL DIAGNOSTIC MODELS FOR LOSS RESERVING	Ben Zehnwrith
35	SEPT 1996	KALMAN FILTERS WITH APPLICATIONS TO LOSS RESERVING	Ben Zehnwrith
36	OCT 1996	RELATIVE REINSURANCE RETENTION LEVELS	David C M Dickson Howard R Waters
37	OCT 1996	SMOOTHNESS CRITERIA FOR MULTI-DIMENSIONAL WHITTAKER GRADUATION	Greg Taylor
38	OCT 1996	GEOGRAPHIC PREMIUM RATING BY WHITTAKER SPATIAL SMOOTHING	Greg Taylor
39	OCT 1996	RISK, CAPITAL AND PROFIT IN INSURANCE	Greg Taylor
40	OCT 1996	SETTING A BONUS-MALUS SCALE IN THE PRESENCE OF OTHER RATING FACTORS	Greg Taylor
41	NOV 1996	CALCULATIONS AND DIAGNOSTICS FOR LINK RATION TECHNIQUES	Ben Zehnwrith Glen Barnett
42	DEC 1996	VIDEO-CONFERENCING IN ACTUARIAL STUDIES – A THREE YEAR CASE STUDY	David M Knox
43	DEC 1996	ALTERNATIVE RETIREMENT INCOME ARRANGEMENTS AND LIFETIME INCOME INEQUALITY: LESSONS FROM AUSTRALIA	Margaret E Atkinson John Creedy David M Knox
44	JAN 1997	AN ANALYSIS OF PENSIONER MORTALITY BY PRE-RETIREMENT INCOME	David M Knox Andrew Tomlin
45	JUL 1997	TECHNICAL ASPECTS OF DOMESTIC LINES PRICING	Greg Taylor
46	AUG 1997	RUIN PROBABILITIES WITH COMPOUNDING ASSETS	David C M Dickson Howard R Waters
47	NOV 1997	ON NUMERICAL EVALUATION OF FINITE TIME RUIN PROBABILITIES	David C M Dickson

No.	Date	Subject	Author
48	NOV 1997	ON THE MOMENTS OF RUIN AND RECOVERY TIMES	Alfredo G Egídio dos Reis
49	JAN 1998	A DECOMPOSITION OF ACTUARIAL SURPLUS AND APPLICATIONS	Daniel Dufresne
50	JAN 1998	PARTICIPATION PROFILES OF AUSTRALIAN WOMEN	M. E. Atkinson Roslyn Cornish
51	MAR 1998	PRICING THE STOCHASTIC VOLATILITY PUT OPTION OF BANKS' CREDIT LINE COMMITMENTS	J.P. Chateau Daniel Dufresne
52	MAR 1998	ON ROBUST ESTIMATION IN BÜHLMANN STRAUB'S CREDIBILITY MODEL	José Garrido Georgios Pitselis
53	MAR 1998	AN ANALYSIS OF THE EQUITY IMPLICATIONS OF RECENT TAXATION CHANGES TO AUSTRALIAN SUPERANNUATION	David M Knox M. E. Atkinson Susan Donath
54	APR 1998	TAX REFORM AND SUPERANNUATION – AN OPPORTUNITY TO BE GRASPED.	David M Knox
55	APR 1998	SUPER BENEFITS? ESTIMATES OF THE RETIREMENT INCOMES THAT AUSTRALIAN WOMEN WILL RECEIVE FROM SUPERANNUATION	Susan Donath
56	APR 1998	A UNIFIED APPROACH TO THE STUDY OF TAIL PROBABILITIES OF COMPOUND DISTRIBUTIONS	Jun Cai José Garrido
57	MAY 1998	THE DE PRIL TRANSFORM OF A COMPOUND R_k DISTRIBUTION	Bjørn Sundt Okechukwu Ekuma
58	MAY 1998	ON MULTIVARIATE PANJER RECURSIONS	Bjørn Sundt
59	MAY 1998	THE MULTIVARIATE DE PRIL TRANSFORM	Bjørn Sundt
60	JUNE 1998	ON ERROR BOUNDS FOR MULTIVARIATE DISTRIBUTIONS	Bjørn Sundt
61	JUNE 1998	THE EQUITY IMPLICATIONS OF CHANGING THE TAX BASIS FOR PENSION FUNDS	M E Atkinson John Creedy David Knox
62	JUNE 1998	ACCELERATED SIMULATION FOR PRICING ASIAN OPTIONS	Felisa J Vázquez-Abad Daniel Dufresne
63	JUNE 1998	AN AFFINE PROPERTY OF THE RECIPROCAL ASIAN OPTION PROCESS	Daniel Dufresne
64	AUG 1998	RUIN PROBLEMS FOR PHASE-TYPE(2) RISK PROCESSES	David C M Dickson Christian Hipp
65	AUG 1998	COMPARISON OF METHODS FOR EVALUATION OF THE n -FOLD CONVOLUTION OF AN ARITHMETIC DISTRIBUTION	Bjørn Sundt David C M Dickson

No.	Date	Subject	Author
66	NOV 1998	COMPARISON OF METHODS FOR EVALUATION OF THE CONVOLUTION OF TWO COMPOUND R_1 DISTRIBUTIONS	David C M Dickson Bjørn Sundt
67	NOV 1998	PENSION FUNDING WITH MOVING AVERAGE RATES OF RETURN	Diane Bédard Daniel Dufresne
68	DEC 1998	MULTI-PERIOD AGGREGATE LOSS DISTRIBUTIONS FOR A LIFE PORTFOLIO	David C M Dickson Howard R Waters
69	FEB 1999	LAGUERRE SERIES FOR ASIAN AND OTHER OPTIONS	Daniel Dufresne
70	MAR 1999	THE DEVELOPMENT OF SOME CHARACTERISTICS FOR EQUITABLE NATIONAL RETIREMENT INCOME SYSTEMS	David Knox Roslyn Cornish
71	APR 1999	A PROPOSAL FOR INTEGRATING AUSTRALIA'S RETIREMENT INCOME POLICY	David Knox
72	NOV 1999	THE STATISTICAL DISTRIBUTION OF INCURRED LOSSES AND ITS EVOLUTION OVER TIME I: NON-PARAMETRIC MODELS	Greg Taylor
73	NOV 1999	THE STATISTICAL DISTRIBUTION OF INCURRED LOSSES AND ITS EVOLUTION OVER TIME II: PARAMETRIC MODELS	Greg Taylor
74	DEC 1999	ON THE VANDERMONDE MATRIX AND ITS ROLE IN MATHEMATICAL FINANCE	Ragnar Norberg
75	DEC 1999	A MARKOV CHAIN FINANCIAL MARKET	Ragnar Norberg
76	MAR 2000	STOCHASTIC PROCESSES: LEARNING THE LANGUAGE	A J G Cairns D C M Dickson A S Macdonald H R Waters M Willder
77	MAR 2000	ON THE TIME TO RUIN FOR ERLANG(2) RISK PROCESSES	David C M Dickson
78	JULY 2000	RISK AND DISCOUNTED LOSS RESERVES	Greg Taylor
79	JULY 2000	STOCHASTIC CONTROL OF FUNDING SYSTEMS	Greg Taylor
80	NOV 2000	MEASURING THE EFFECTS OF REINSURANCE BY THE ADJUSTMENT COEFFICIENT IN THE SPARRE ANDERSON MODEL	Maria de Lourdes Centeno
81	NOV 2000	THE STATISTICAL DISTRIBUTION OF INCURRED LOSSES AND ITS EVOLUTION OVER TIME III: DYNAMIC MODELS	Greg Taylor

No.	Date	Subject	Author
82	DEC 2000	OPTIMAL INVESTMENT FOR INVESTORS WITH STATE DEPENDENT INCOME, AND FOR INSURERS	Christian Hipp
83	DEC 2000	HEDGING IN INCOMPLETE MARKETS AND OPTIMAL CONTROL	Christian Hipp Michael Taksar
84	FEB 2001	DISCRETE TIME RISK MODELS UNDER STOCHASTIC FORCES OF INTEREST	Jun Cai
85	FEB 2001	MODERN LANDMARKS IN ACTUARIAL SCIENCE Inaugural Professorial Address	David C M Dickson
86	JUNE 2001	LUNDBERG INEQUALITIES FOR RENEWAL EQUATIONS	Gordon E Willmot Jun Cai X Sheldon Lin
87	SEPTEMBER 2001	VOLATILITY, BETA AND RETURN WAS THERE EVER A MEANINGFUL RELATIONSHIP?	Richard Fitzherbert
88	NOVEMBER 2001	EXPLICIT, FINITE TIME RUIN PROBABILITIES FOR DISCRETE, DEPENDENT CLAIMS	Zvetan G Ignatov Vladimir K Kaishev Rossen S Krachunov
89	NOVEMBER 2001	ON THE DISTRIBUTION OF THE DEFICIT AT RUIN WHEN CLAIMS ARE PHASE-TYPE	Steve Drekcic David C M Dickson David A Stanford Gordon E Willmot
90	NOVEMBER 2001	THE INTEGRATED SQUARE-ROOT PROCESS	Daniel Dufresne