


Two radio relics discovered in the galaxy cluster Abell 168

March 28 2018, by Tomasz Nowakowski



GMRT images of Abell 168. Credit: Dwarakanath et al., 2018.

Indian astronomers have detected two radio relics in the nearby merging galaxy cluster known as Abell 168. The discovery, which could improve our understanding of merging processes in galaxy clusters, was detailed March 16 in a paper published on the arXiv pre-print repository.

Radio relics are diffuse, elongated radio sources of synchrotron origin. They occur in the form of spectacular single or double symmetric arcs at the peripheries of [galaxy clusters](#). Astronomers are especially interested in searching for such sources in merging galaxy clusters as the number of radio relics associated with merger shocks is still small.

Abell 168 is a nearby low-mass merging galaxy cluster at a redshift of 0.045 with a mass of about 152 trillion solar masses. First observations of radio emission associated with this cluster date back to 2011. Now, a team of researchers led by K. S. Dwarakanath of the Raman Research Institute in Bangalore, India, reports the finding of two radio relics in the Abell 168 galaxy cluster. The discovery was made using the Giant Metrewave Radio Telescope (GMRT) and the Karl G. Jansky Very Large Array (VLA).

"We report the discovery of twin radio relics in the outskirts of the low-mass merging galaxy cluster Abell 168 (redshift=0.045)," the researchers wrote in the paper.

GMRT and VLA observations revealed that one of the relics is elongated, thin and located toward the northern periphery of Abell 168 at a distance of about 3 million light-years from the cluster center. This relic has a linear extent of approximately 2.6 million light-years and a

projected width of about 260,000 light-years.

The second newly found radio relic is ring-shaped and has a size of around 700,000 light-years. This radio source is located near the inner edge of the elongated relic, some two million light-years from the center of Abell 168. Moreover, the researchers found that the ring-shaped relic (source A) has a steeper spectral index of about -1.74 in the frequency range between 100 and 600 MHz and appears to be connected to the elongated relic (source B). They assume that the ring-shaped relic is old plasma revived due to adiabatic compression by the outgoing shock that produced the elongated relic.

"The orientation of source B, the projected distance between sources A and B, and the radiative lifetime of source A are consistent with such a scenario," the paper reads.

In concluding remarks, the authors noted that their study, besides identifying two new radio relics in Abell 168, provides an important information about the galaxy [cluster](#) itself. They emphasized that the discovery makes Abell 168 the lowest mass system so far known to have radio relics of merger shock origin.

More information: Twin radio relics in a near-by low-mass galaxy cluster Abell 168, arXiv:1803.06129 [astro-ph.CO]
arxiv.org/abs/1803.06129

Abstract

We report the discovery of twin radio relics in the outskirts of the low-mass merging galaxy cluster Abell 168 (redshift=0.045). One of the relics is elongated with a linear extent ~ 800 kpc, a projected width of ~ 80 kpc and is located ~ 900 kpc toward the north of the cluster center, oriented roughly perpendicular to the major axis of the X-ray emission. The second relic is ring-shaped with a size ~ 220 kpc and is located near

the inner edge of the elongated relic at a distance of ~ 600 kpc from the cluster center. These radio sources were imaged at 323 and 608 MHz with the Giant Meterwave Radio Telescope and at 1520 MHz with the Karl G Jansky Very Large Array (VLA). The elongated relic was detected at all the frequencies with a radio power at 1.4 GHz of $1.38 \pm 0.14 \times 10^{23} \text{ W Hz}^{-1}$ having a power law in the frequency range 70 - 1500 MHz ($S \propto \nu^\alpha, \alpha = -1.1 \pm 0.04$). This radio power is in good agreement with that expected from the known empirical relation between the radio powers of relics and the host cluster masses. This is the lowest mass ($M_{500} = 1.24 \times 10^{14} \text{ Mo}$) cluster in which relics due to merger shocks are detected. The ring-shaped relic has a steeper spectral index (α) of -1.74 ± 0.29 in the frequency range 100 - 600 MHz. We propose this relic to be an old plasma revived due to adiabatic compression by the outgoing shock which produced the elongated relic.

© 2018 Phys.org

Citation: Two radio relics discovered in the galaxy cluster Abell 168 (2018, March 28) retrieved 12 September 2024 from <https://phys.org/news/2018-03-radio-relics-galaxy-cluster-abell.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.