


# New dataset reveals accelerated global soil phosphorus release at higher temperatures

July 15 2024, by Zhang Nannan



A schematic diagram for soil phosphorus cycling under various weathering stages. Credit: *Science Advances* (2024). DOI: 10.1126/sciadv.adm7773

A study [published](#) in *Science Advances* shows that phosphorus (P) release from soils is enhanced at higher mean annual temperatures (MAT). This finding is based on a new compilation of data on global surface soil temperatures and phosphorus content.

Phosphorus released from soils via [chemical weathering](#) (referred to as P weathering) plays a critical role in the global cycling of key elements and influences the size of the Earth's biosphere. While it has been theoretically proposed that [global climate](#) significantly affects P weathering, direct empirical evidence on a global scale has been lacking

until now.

In this study, Profs. Guo Licheng, Zhao Mingyu, Xiong Shangfa, and Yang Shiling from the Institute of Geology and Geophysics of the Chinese Academy of Sciences (CAS), together with their collaborators, compiled a geochemical dataset of global surface soils to investigate the relationship between climate and P weathering.

The dataset shows that temperature is the primary regulator of P mobility. It shows reduced P retention in soils in [warmer climates](#) (> 12°C) and with high silicate weathering intensities, characterized by the near-complete leaching of Na<sup>+</sup>, Ca<sup>2+</sup>, K<sup>+</sup> from fresh regolith.

Further analysis indicates that lower [soil](#) pH in high weathering intensity environments promotes the removal of primary apatite and the dissolution of Al, Fe, Ca phosphates. In addition, in such environments, a higher kaolinite/illite ratio results in a decreased P adsorption capacity within clay minerals.

The researchers also calculated the relationship between modern global MAT and P weathering flux based on the observed relationship between MAT and P content, as well as the modern latitudinal distributions of temperature and land area. The model results show a rapid increase in P weathering flux within the global MAT range of 20°C to 23°C.

Considering the positive effects of nutrient supply on primary productivity and organic matter burial, this study suggests that enhanced P weathering flux in warm climates is a critical component of the Earth's natural thermostat and was likely the cause of oceanic anoxia during past climate warming events.

"A further implication of this study is that the potential acceleration of P loss from soils due to anthropogenic climate warming may pose threats

to [agricultural production](#), terrestrial and [marine ecosystems](#), and alter marine redox landscapes," said Prof. Zhao, corresponding author of the study.

**More information:** Licheng Guo et al, Acceleration of phosphorus weathering under warm climates, *Science Advances* (2024). [DOI: 10.1126/sciadv.adm7773](#)

Provided by Chinese Academy of Sciences

Citation: New dataset reveals accelerated global soil phosphorus release at higher temperatures (2024, July 15) retrieved 20 August 2024 from <https://phys.org/news/2024-07-dataset-reveals-global-soil-phosphorus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.