



E-ISSN: 2320-7078
P-ISSN: 2349-6800
JEZS 2017; 5(3): 365-369
© 2017 JEZS
Received: 23-03-2017
Accepted: 24-04-2017

AK Patil
Department of Animal
Nutrition, College of Veterinary
Science & A.H., Jabalpur,
Madhya Pradesh, India

RPS Baghel
Department of Animal
Nutrition, College of Veterinary
Science & A.H., Jabalpur,
Madhya Pradesh, India

S Nayak
Department of Animal
Nutrition, College of Veterinary
Science & A.H., Jabalpur,
Madhya Pradesh, India

CD Malapure
Department of Animal
Nutrition, College of Veterinary
Science & A.H., Jabalpur,
Madhya Pradesh, India

K Govil
Department of Animal
Nutrition, College of Veterinary
Science & A.H., Jabalpur,
Madhya Pradesh, India

D Kumar
Department of Veterinary
Science & A.H., COA, JNKVV,
Tikamgarh- 472001, Madhya
Pradesh, India

Pavan Kumar Yadav
Faculty of Veterinary and
Animal Sciences, IAS, Banaras
Hindu University, Varanasi,
Uttar Pradesh, India

Correspondence
Pavan Kumar Yadav
Assistant Professor (Veterinary
Biochemistry), Department of
Physiology and Biochemistry,
Faculty of Veterinary and
Animal Sciences, IAS, Banaras
Hindu University, Varanasi-
221005, Uttar Pradesh, India

Cumin (*Cuminum cyminum*): As a Feed Additive for Livestock

AK Patil, RPS Baghel, S Nayak, CD Malapure, K Govil, D Kumar and Pavan Kumar Yadav

Abstract

Cumin is an annual herb and has been used since ancient times as medicine and spices in food. Supplementation of cumin could either influence the feeding pattern or growth of favorable microorganisms in the rumen or stimulate the secretion of various digestive enzymes, which in turn may improve the efficiency of nutrients utilization or stimulate the milk secretory tissues in mammary glands resulting in improved milk production and reproductive performance of dairy animals. It has various pharmacological effects; recently the use of cumin has gained popularity because of herbal movement initiated by naturopaths, yog gurus, alternative medicine promoters and feed additives. Animal nutritionist is trying to exploit the potential use of cumin as a growth promoter, efficiency of nutrient utilization and mitigation of greenhouse gas emission. This review highlights the potential use of cumin as feed additive to increase production efficiency for effective animal production and reproduction.

Keywords: Cumin, feed additive, livestock, pharmacological effects, seed spices

1. Introduction

Cumin (*Cuminum cyminum* Linn.) is an important commercial seed spice belonging to the family umbelliferae and known for its aroma, medicinal and therapeutic properties [1]. The plant is native to the Mediterranean region where it is cultivated extensively. It is commonly known as zeera/jeera in Hindi, is an annual herb and has been used since ancient times as medicine and spices in food. It is considered to be one of the most important spices and ranks second to black pepper [2]. India is the world's largest producer and consumer of cumin [3]. Many value-added products of cumin seeds like oleoresins of cumin seeds and cumin oil are also exported from India. This spice should not be confused with sweet cumin, which is a common name for anise (*Pimpinella anisum*) [4]. *Nigella sativa*, which is also frequently referred to as black cumin, is not related to cumin. World widely therapeutic uses of cumin have been experimentally validated. In Ayurveda cumin seeds are notably considered carminative, eupeptic, antispasmodic, astringent and are used for treating mild digestive disorders, diarrhea, dyspepsia, flatulence, colic, abdominal distension, edema, broncho-pulmonary disorders, puerperal disorders, analgesic and as a cough remedy [5]. Cumin has various actions like enhancing vision, strength and lactation [6]. Cumin seeds have been reportedly used for traditional treatment of toothache, dyspepsia, epilepsy and jaundice [7]. It has various pharmacological effects such as anti-diabetic, immunologic, anti-tumour and antimicrobial activities [8]. Essential oil of cumin acts as powerful external or internal antiseptic, an analgesic, anti-inflammatory, hemolytic, anti-enzymatic action, sedative, stimulants and stomachic's [9]. Iranian traditional medicine consider cumin as stimulant, carminative and astringent and its therapeutic effects have been described on gastrointestinal, gynecological and respiratory disorders and also for the treatment of toothache, diarrhea and epilepsy [10]. In traditional medicine of Tunisia, cumin is considered abortive, galactagogue, antiseptic, antihypertensive herb, while in Italy; it is used as bitter tonic, carminative and purgative [11]. Therefore, it is clear that cumin as feed additives have many beneficial effects due to its aforesaid properties. Research has showed that the use of chemical additives especially antibiotics in animal nutrition may result in the accumulation of chemical residues in animal products [12]. Therefore, the use of organic feed additives in animal nutrition has gained more attention during recent years due to concerns about food safety and human health issue.

2. Composition

Cumin seed contains several nutrients including moisture (7%), volatile oil (3-4%), protein (12%), total ash (10%), fiber (11%), carbohydrate (33%), starch (11%) and fat (15%) [13]. Its nutrients composition varies according to the region and climate condition where it is grown. Typical cumin flavor is due to the volatile oil present in the seeds in the range of 3-4% depending on the variety and the origin of the cultivation. The cumin seeds contain 14.5% total lipids on dry weight basis which is consisted of 84.8% neutral lipid, 10% glycolipids and 5.1% phospholipids. Fatty acid composition of cumin includes both saturated and unsaturated fatty acids mainly (70% of total) oleic, petroselinic and linoleic acids [14]. Cumin essential oil contains several compounds namely cumin aldehyde, β -pinene, p-cymene and γ -terpinene [15]. Good amount of phenolic compounds show considerable radical scavenging carotene/ linoleic acid chelating and reducing power activities [16].

3. Effect of cumin supplementation on the performance of animals

3.1 Effect on nutrient utilization and methane mitigation

Livestock contributes about 18% to the global anthropogenic greenhouse gas (GHG) emissions, accounting for about 37% of the total anthropogenic methane and 65% of global anthropogenic nitrous oxide [17]. Ruminant livestock such as cattle can produce up to 200 L of methane per day which is regarded as a significant threat to the environment. Now a day's more emphasis is given for reducing methane from ruminants because eructation of methane from ruminants may contribute to the climatic change and global warming [18]. Moreover, ruminant animals lose about 2-12% of their dietary gross energy as methane which affects the nutrient efficiency and significantly contributes to the greenhouse gas emissions. Spices that have long been safely used for human consumption could be tested as alternatives to reduce enteric methane from livestock. Some spices are rich in tannins or saponins or polyphenolics and others are high in fatty acids [19]. Several researchers [20, 21] have used plant extracts to manipulate rumen fermentation for improving nutrient utilization and minimizing methane emission.

Number of researcher has proved that the inclusion of cumin increase digestibility of nutrients and enhance nutrient utilization additionally, it has potential to reduce methane emission from ruminants [22, 23]. Contrary, Miri *et al.* [24] observed that the DMI and milk composition were unaffected after addition of cumin seed in the ration, indicating no negative effect of cumin seed on palatability of the ration.

Additions of cumin have potential to reduced methane to the extent of 22% in buffered rumen fluid, *in vitro* observed that there was 11.8% reduction in methane emission of goats when diet supplemented with cumin seed extract [25]. This effect might be due to inhibitory effect of cumin seed extract on ruminal microbial biomass. Suppression of growth of ruminal Gram-positive H₂ producing bacteria can reduce the amount of available hydrogen for biohydrogenation as well as methanogenesis. The loss of protozoa would be expected to impact positively on the availability of CLA to host animal and negatively on methane emission [26].

Supplementing broiler diet with cumin oil has significantly increased body weight as compared with control [27]. Ali *et al.* [28] reported that negative effects of heat stress can be alleviated and growth performance can be improved by adding 2% cumin in diet of heat stressed broiler. Galib and Al-Kassi [29] reported that adding cumin and turmeric mixture

at levels of 0.75% and 1% in the diets improved body weight gain, feed intake and feed conversion ratio in broilers, which could be due to role of cumin as a stimulant, carminative, digestion, antimicrobial properties and the prevention of gastric toxicity. Other researchers proved that there is an increase in body weight, feed conversion ratio with decreasing hematological values of some important blood parameters using 2% of cumin in broiler diets [30].

3.2 Effect on milk quantity and quality

Sustainable dairy farming necessitates new approaches to improve milk production efficiency. The need for more efficient production systems is growing due to the ongoing increases in the costs of production particularly those of feedstuff in developing countries such as India. Cumin is a constituent of herb Payapro a known galactagogue. Bhatt *et al.* [31] also reported galactapoiesis property. The addition of cumin seed extract (1.27% of DM) significantly ($P < 0.05$) enhanced milk production by 13% in supplemented goat in comparison to control group [24]. Cumin seed extract (CSE) supplementation had no any adverse effect on milk fat, protein and lactose percentage, however, C18:3-n3, C18:2n-6c, C18:1 trans-11, monounsaturated fatty acids, polyunsaturated fatty acids (PUFA) and ratio of polyunsaturated to saturated fatty acids content of milk increased significantly ($P < 0.05$) when additive was used [32]. The cis-9 trans-11 conjugated linoleic acid (CLA) significantly ($P < 0.01$) increased by 20% in goat milk receiving the CSE supplemented diet. Delta-9-desaturase index in CSE added groups was significantly ($P < 0.01$) higher in comparison to control. In addition, goats which were fed with 1g/L CSE produced milk with higher recovery of linoleic acid (LA) and linolenic acid (LNA) [32]. Polyphenols (tannins and non-tannins) have ability to affect fatty acid metabolism at different steps of ruminal biohydrogenation [33]. Bettaieb *et al.* [34] observed that cumin contains an unusual fatty acid like petroselinic acid (C18:1n-12)) and part of the increase in the concentration of LA in goat milk fed CSE might be attributed to the presence of petroselinic acid, that has been shown to inhibit conversion of LA to arachidonic acid [35]. Likewise, Ghafari *et al.* [36] observed that the supplementation of cumin seed @ 0, 100, 200 and 300 g/cow/d have potential to enhanced ($P \leq 0.05$) milk yield curvilinearly with level of cumin seed (average 47.9, 52.5, 55.1 and 53.6 kg/d for the four levels, respectively). The yield of milk components was similar to milk yield ($P \leq 0.05$) except for fat and 4% fat-corrected milk yields, which were not significantly affected by the treatments.

3.3 Effect on antioxidant status

Free radicals are highly reactive species having unpaired electrons in their outermost shell [37]. Higher production of free radicals in the body can causes oxidative stress that ultimately leads to oxidative damage to important biomolecules, leading to many chronic diseases [38]. To cope up with these radicals the living system has antioxidant enzymes defense system or animal may takes antioxidants through diet as vitamins and minerals. Number of studies conducted in India showed that cumin oils exhibited high antioxidant activity due to flavonoids particularly apigenin and luteolin of cumin seeds [39]. Gangandeeep [40] reported that supplementation of cumin seed (2.5 and 5% of diet) in mice tended to increase superoxide dismutase, catalase and reduced glutathione however, the activities of glutathione peroxidase and glutathione reductase remained unaltered by both doses of

cumin. Similarly, cuminaldehyde has been demonstrated to scavenge the superoxide anion [41]. Further it confirmed by Juhainmi and Ghafoor [42] that the supplementation of cumin seed extracts has higher DPPH radical scavenging activities in comparison to unsupplemented animals and the antioxidant activity of cumin seed extracts ranged from 8.25 to 11.24 mg/mL. They stated that the antioxidative potential increased with increase of phenolic compounds content of cumin. Food materials rich in bioactive compounds with higher free radical scavenging abilities are protective against certain types of cancer and may also reduce the risk of cardiovascular and cerebrovascular disorders in human being [43].

3.4. Antimicrobial activity

The major compounds in all cumin oils were the monoterpenes beta-pinene, p-cymene, gamma-terpinene, terpenoid aldehydes cuminic aldehyde and isomeric menthadien carboxaldehydes [44]. Cumin essential oil showed antibacterial activity comparable with standard antibiotics against common pathogens. Cuminaldehyde demonstrated antimicrobial and antifungal properties against *E. coli*, *P. chrysogenum*, *A. flavus* and *A. niger* [45]. Similarly, Jirovets *et al.* [46] also reported anti-microbial activity of cumin essential oil against molds (*A. niger*), gram positive bacteria (*B. subtilis*, *S. epidermidis*), gram negative bacteria (*E. coli*) and yeast (*S. cerevisiae* and *C. albicans*). Jazani *et al.* [47] indicated the potential use of cumin essential oil for the control of some diseases caused by *P. aeruginosa* infections. Cumin oil and cuminaldehyde have been reported to exhibit strong larvicidal and antibacterial activity. At *in vitro* concentrations of 300 or 600 ppm, cumin oil inhibited the growth of *Lactobacillus plantarum* [48]. Antifungal activity of cumin oil is recorded against soil, food, animal and human pathogens, including dermatophytes, *Vibrio* spp., yeasts, aflatoxins and mycotoxin producers [49, 50, 51].

3.5 Immunomodulatory effect

Oral treatment with cumin caused modulation of T-lymphocyte's expression in a dose dependent manner in normal and immune suppressed animals. It stimulated the T-cell's (CD₄ and CD₈) and Th-1 cytokine's expression in normal and cyclosporine-A induced immune suppressed mice. In stress induced immune suppressed animals the active compound of cumin countered the depleted T lymphocytes, decreased the elevated corticosterone levels and size of adrenal glands and increased the weight of thymus and spleen [52].

3.6 Estrogenic/anti-osteoporotic effect

Cumin seeds content phytoestrogens which is responsible for estrogenic and anti-osteoporotic property [53]. Administration of cumin extract in animals resulted in reduced urinary calcium excretion and increased calcium content and bone strength. Bone and ash densities along with improved microarchitecture with no adverse effects were reported after supplementation of cumin in animal [54].

3.7 Anticarcinogenic/antimutagenic

Gangandee *et al.* [40] observed that cumin seeds were able to inhibit the induction of gastric squamous cell carcinomas in mice. Similarly, in rats fed with cumin has a protective effect against induced colonic cancer, decreased beta-glucuronidase and mucinase activity have been demonstrated [55]. Dietary cumin inhibited benzopyrene-induced forestomach tumorigenesis, 3-methylcholanthrene induced uterine cervix

tumorigenesis and 3-methyl-4-dimethylaminoazobenzene induced hepatomas in mice [56, 57]. This was attributed to the ability of cumin in modulating carcinogen metabolism via carcinogen/xenobiotic metabolizing phase I and phase II enzymes. Activities of cytochrome (CYP) P-450 reductase and CYP b5 reductase were augmented whereas phase II enzymes GST and DT-diaphorase were increased [58].

3.8 Antidiabetic effect

The antidiabetic effects of cumin are well documented [59]. In a glucose tolerance test in rabbits, cumin significantly increased the area under the glucose tolerance curve and hyperglycemic peak [60]. Diet containing cumin powder (1.25%) was found to be remarkably beneficial in streptozotocin induced diabetic rats as indicated by reduction in hyperglycemia and glucosuria, improved body weights, lowered blood urea level and reduced excretions of urea and creatinine by diabetic animals [61]. Cumin at (0.5% g) per kg body weight orally administered in rats with induced diabetes is reported reduce blood glucose levels [62]. It might be due to inhibition of aldose reductase and alpha-glucosidase [63].

4. Conclusion

Cumin is one of the commercially important seed spices and forms one of the ingredients of many spice mixes which are consumed in our daily diet. *Cuminum cyminum* contained alkaloid, coumarin, anthraquinone, flavonoid, glycoside, protein, resin, saponin, tannin and steroid. Its fatty acid composition and cumin oleoresin has rich in unsaturated fatty acids mainly linoleic acid which is known to possess health benefits. Flavanoids like apigenin, luteolin and glycosides present in cumin are reported to be responsible for many of the biological activities of cumin. Cumin acts as antimicrobial, insecticidal, anti-inflammatory, analgesic, antioxidant, anticancer, antidiabetic, antiplatelet aggregation, hypotensive, bronchodilatory, immunological, anti-amyloidogenic and anti-osteoporotic effects etc. From the review, it is concluded that cumin as feed additives is included in small quantities in the diet to improve nutrient utilization and production efficiency without any adverse impact on animals.

5. Acknowledgement

We declare that we have no conflicts of interest.

6. References

1. Sowbhagya HB. Chemistry, Technology and Nutraceutical Functions of Cumin *Cuminum cyminum* L.: An Overview. Critical Reviews in Food Science and Nutrition. 2013; 53(1):1-10.
2. Kitchen dictionary. food.com. 2014. <http://www.food.com/library/cumin-20>.
3. Spices Board, India, 2016. www.indianspices.com.
4. Gilligan NP. The palliation of nausea in hospice and palliative care patients with essential oils of *Pimpinella anisum* (aniseed), *Foeniculum vulgare* var. *dulce* (sweet fennel), *Anthemis nobilis* (Roman chamomile) and *Mentha x piperita* (peppermint). International Journal of Aromatherapy. 2005; 15(4):163-167.
5. Nadkarni KM, Nadkarni AK. Indian Material Medica. Popular Prakashan Pvt. Ltd, Mumbai, India, 1976.
6. Rathore SS, Saxena SN, Singh B. Potential health benefits of major seed spices. Indian Society for Seeds and Spices. 2013; 3:1-12.
7. Nostro A, Cellini L, Di Bartolomeo S, Di Campli E,

- Grande R, Cannatelli MA. Antibacterial effect of plant extracts against *Helicobacter pylori*. *Phytotherapy Research*. 2005; 19:198-202.
8. Senevirathne M, Kim S, Siriwardhana N, Ha J, Lee K, Jeon Y. Antioxidant potential of *Ecklonia cava* on reactive oxygen species scavenging metal chelating, reducing power and lipid peroxidation inhibition. *International Journal of Food Science & Technology*. 2006; 12:27-38.
 9. Guenther E. *The Essential Oils*. 3rd Ed. D. Van Nostrand, New York, 1950.
 10. Zargary A. 5th ed. Tehran: Tehran University Publications. *Medicinal Plants in Persian*, 2001.
 11. Leporatti ML, Ghedira K. Comparative analysis of medicinal plants used in traditional medicine in Italy and Tunisia. *Journal of Ethnobiology and Ethnomedicine*. 2009; 5:31-39.
 12. Amir Davar Foroozandeh, Mojtaba Ghaffari. The effects of feeding different levels of *Cuminum cyminum* L. on milk yield and composition and some blood metabolites of Holstein dairy cows. *Proceedings of the IRES 4th International Conference*, Kuala Lumpur, Malaysia, 2015. ISBN: 978-93-85465-56-7.
 13. Lewis YS. *Spices and herbs for the Food Industry*, England: Food Trade press. 1984, 121-122.
 14. Hemavathy J, Prabhakar JV. Lipid composition of cumin *Cuminum cyminum* L. seeds. *Journal of Food Science*. 1988; 53(5):1578-79.
 15. Heri YJ, Wen T, Ming Z, Ninghua D. Determination of chemical components of volatile oil from *Cuminum cyminum* L. by gas chromatography mass spectrometry. *Peop. Rep. China*. 2003; 41:4000.
 16. Rebey IB, Jabri-Karoui I, Hamrouni-Sellami I, Bourgou S, Limam F, Marzouk B. Effect of drought on the biochemical composition and antioxidant activities of cumin *Cuminum cyminum* L. seeds. *Industrial Crops and Products*. 2012; 36:238-245.
 17. FAOSTAT. *Food and Agriculture Organization of the United Nations*, 2006.
 18. Hansen J. A slippery slope: How much global warming constitutes “dangerous anthropogenic interference. *Climate change*. 2005; 68(333):269-279. Doi: 10.1007/s10584-005-4135-0.
 19. Rai N, Yadav S, Verma AK, Tiwari L, Sharma RK. A monographic profile on quality specifications for a herbal drug and spice of commerce- *Cuminum cyminum* L. *International Journal of Advanced Herbal Science and Technology*. 2012; 1(1):1-12.
 20. Benchaar C, McAllister TA, Chouinard PY. Digestion, ruminal fermentation, ciliate protozoal populations and milk production from dairy cows fed cinnamaldehyde, quebracho condensed tannin, or *Yucca schidigera* saponin Extracts. *Journal of Dairy Science*. 2008; 91:4765-77.
 21. Patra AK, Kamra DN, Agarwal N. Effect of plant extracts on *in vitro* methanogenesis: enzyme activities and fermentation of feed in rumen liquor of buffalo. *Anim. Animal Feed Science and Technology*. 2006; 128:276-291.
 22. Khan MMH, Chaudhry AS. Chemical composition of selected forages and spices and the effect of these spices on *in vitro* rumen degradability of some forages. *Asian-Australasian Journal of Animal Sciences*. 2010; 23:889-900.
 23. Kilic U, Boga M, Gorgulu MS, Ahan Z. The effects of different compounds in some essential oils on *in vitro* gas production. *The Journal of Animal and Feed Sciences*. 2011; 20:626-636.
 24. Miri H, Tyagi V, Ebrahimi AK, Hadi Mohini S. Effect of cumin *Cuminum cyminum* seed extract on milk fatty acid profile and methane emission in lactating goat. 2013; 113:66-72.
 25. Chaudhry AS, Khan MMH. Impacts of different spices on *in vitro* rumen dry matter disappearance, fermentation and methane of wheat or ryegrass hay based substrates. *Livestock Science*. 2012; 146:84-90.
 26. Lourenco M, Ramos-Morales E, Wallace RJ. The role of microbes in rumen lipolysis and biohydrogenation and their manipulation. *Animal*. 2010; 4:1008-1023.
 27. Al-Anbari EH, Abbas, AA, Al-Samarai FR, Al-Shamire JS, Al-Zaidi FH. Effect of using cumin oil *Cuminum cyminum* as feed additives on profile analysis and growth curve of broiler GJBB. 2013; 2:326-330.
 28. Ali MN, Qota NMA, Hassan RA. Recovery from adverse effects of heat stress on slow growing chicks using natural antioxidant with or without sulfate, *International Journal of Poultry Science*. 2010; 9:109-117.
 29. Galib AM, Al-Kassi. Effect of feeding cumin *Cuminum cyminum* on the performance and some blood traits of broiler chicks. *Pakistan Journal of Nutrition*. 2010; 9:72-75.
 30. Ibrahim IA, El Badwi SMA, Bakhiet AO, Gadir WS Abdel, Adam SEI. A 9-week feeding study of *Cuminum cyminum*. *Journal of Pharmacology and Toxicology*. 2007; 2:666-671.
 31. Bhatt N, Singh M, Ali A. Effect of feeding herbal preparations on milk yield and rumen parameters in lactating crossbred cows. *International Journal of Agriculture and Biology*. 2009; 11:721-726.
 32. Heidarian M, Vahideh, Ebrahimi, Seyed H, Kumar T, Amrish. The effect of cumin *Cuminum cyminum* seed extract on the inhibition of PUFA biohydrogenation in the rumen of lactating goats via changes in the activity of rumen bacteria and linoleate isomerase enzyme. *Small ruminant research*. 2015; 125:56-63.
 33. Jayanegara A, Kreuzer M, Leiber F. Ruminal disappearance of polyunsaturated fatty acids and appearance of biohydrogenation products when incubating linseed oil with alpine forage plant species. *Livestock Science*. 2012; 147:104-112.
 34. Bettaieb I, Bourgou S, Sriti J, Msaada K, Limam F, Marzouk B. Essential oils and fatty acids composition of Tunisian and Indian cumin *Cuminum cyminum* L. seeds: a comparative study. *Journal of the Science of Food and Agriculture*. 2011; 91:2100-2107.
 35. Weber N, Richter KD, Schulte E, Mukherjee KD. Petroselinic acid from dietary triacylglycerols reduces the concentration of arachidonic acid in tissue lipids of rats. *Journal of Nutrition*. 1995; 125:1563-1568.
 36. Ghafaria M, Foroozandeh Shahrakia AD, Nasrollahib SM, Aminib HR, Beauchemin KA. Cumin seed improves nutrient intake and milk production by dairy cows. *Animal Feed Science and Technology*. 2015; 210:276-280.
 37. Hamid AA, Aiyelaagbe OO, Usman LA, Ameen OM, Lawal A. Antioxidants: Its medicinal and pharmacological applications. *African Journal of Pure and Applied Chemistry*. 2010; 4(8):142-151.
 38. Poulson HE, Prieme H, Loft S. Role of oxidative DNA

- damage in cancer initiation and promotion. European Journal of Cancer Prevention. 1998; 7:9-16.
39. Leung AY. Encyclopedia of Common Natural Ingredients used in Foods, Drugs and Cosmetics, John Wiley, Hoboken NJ, 1980.
 40. Gagandeep S, Dhanalakshmi S, Mendiz E, Rao AR, Kale RK. Chemopreventive effects of *Cuminum cyminum* in chemically induced forestomach and uterine cervix tumours in murine model systems. Nutrition and Cancer. 2003; 47:171-80.
 41. Krishnakantha TP, Lokesh BR. Scavenging of superoxide anions by spice principles. Indian Journal of Biochemistry and Biophysics. 1993; 30:133-134.
 42. Juhaimi AL, Ghafoor K. Extraction optimization and *in vitro* antioxidant properties of phenolic compounds from Cumin *Cuminum cyminum* L. seed. International Food Research Journal. 2013; 20:1669-1675.
 43. Miraliakbari H, Shahidi F. Antioxidant activity of minor components of tree nut oils. Food Chemistry. 2008; 111:421-427.
 44. Wanner J, Bail S, Jirovets L, Buchbauer G, Schmidt E, Gochev V, *et al.* Chemical composition and antimicrobial activity of cumin oil *Cuminum cyminum*, Apiaceae. Natural Product Communications. 2010; 5(9):1355-1358.
 45. Singh G, Upadhyay RK. Essential oils: a potent source of natural pesticides. Journal of Scientific and Industrial Research. 1993; 52:676-683.
 46. Jirovets L, Buchbauer G, Stoyanova AS, Georgiev EV, Damianoga ST. Composition, quality control and antimicrobial activity of the essential oil of Cumin *Cuminum cyminum* L. seeds from Bulgaria that had been stored for up to 36 years. International Journal of Food Science and Technology. 2005; 40:305.
 47. Jazani NH, Zartoshti M, Shahabi S. Antibacterial Effects of Iranian *Cuminum cyminum* Essential Oil on Burn of *Pseudomonas aeruginosa*. International Journal of Pharmacology. 2008; 4:157-159.
 48. Kivanc M, Akgul A, Dogan A. Inhibitory and stimulatory effects of cumin, oregano and their essential oils on growth and acid production of *Lactobacillus planatarum* and *Leuconstoc mesenteroides*. International Journal of Food Microbiology. 1991; 13:81-85.
 49. De Martino L, De Feo V, Fratianni F, Nazzaro F. Chemistry, antioxidant, antibacterial and antifungal activities of volatile oils and their components. Natural Product Communications. 2009; 4:1741-50.
 50. Hajlaoui H, Mighri H, Noumi E, Snoussi M, Trabelsi N, Ksouri R. Chemical composition and biological activities of Tunisian *Cuminum cyminum* L. essential oil: A high effectiveness against *Vibrio* spp. strains. Food and Chemical Toxicology. 2010; 48:2186-92.
 51. Mehdi RA, Masoomch SG, Mohammad BR, Kamkar J, Soheil A, Reza S. Chemical composition and antiaflatoxicogenic activity of *Carum carvi* L., *Thymus vulgaris* and *Citrus aurantifolia* essential oils. Food Control. 2009; 20:1018-24.
 52. Chauhan P, Satti NK, Suri KA, Amina M, Bani S. Stimulatory effects of *Cuminum cyminum* and flavonoid glycoside on cyclosporine-A and restraint stress induced immune-suppression in swiss albino mice. Chemico-Biological Interactions. 2010; 185:66-72.
 53. Malini T, Vanithakumari G. Estrogenic activity of *Cuminum cyminum* in rats. Indian Journal of Experimental Biology. 1987; 25:442-4.
 54. Shirke SS, Jadhav SR, Jagtap AG. Methanolic extract of *Cuminum cyminum* inhibits ovariectomy-induced bone loss in rats. Experimental Biology and Medicine. 2008; 233:1403-10.
 55. Nalini N, Sabitha K, Vishwanathan P, Menon VP. Influence of spices on the bacterial (enzyme) activity in experimental colon cancer. Journal of Ethnopharmacology. 1998; 62:15-24.
 56. Kamaleeshwari M, Deeptha K, Sengottuvelan M, Nalini N. Effect of dietary caraway on aberrant crypt foci development, fecal steroids, and intestinal alkaline phosphatase activities in 1,2-dimethylhydrazine-induced colon carcinogenesis. Toxicology and Applied Pharmacology. 2006a; 214:290-6.
 57. Deeptha K, Kamaleeshwari M, Sengottuvelan M, Nalini N. Dose dependent inhibitory effect of dietary caraway on 1, 2-dimethylhydrazine induced colonic aberrant crypt foci and bacterial enzyme activity in rats. Investigational New Drugs. 2006; 24:479-88.
 58. Aruna K, Sivaramakrishnan VM. Anticarcinogenic effects of some Indian plant products. Food and Chemical Toxicology. 1992; 30:953-956.
 59. Srinivasan K. Plant foods in the management of diabetes mellitus: Spices as beneficial antidiabetic food adjuncts. Int. International Journal of Food Sciences and Nutrition. 2005; 56:399-414.
 60. Roman-Ramos R, Flores-Saenz JL, Alarcon-Aguilar FJ. Anti-hyperglycemic effect of some edible plants. Journal of Ethnopharmacology. 1995; 48:25-32.
 61. Willatgamuwa SA, Platell K, Saraswathi G, Srinivasan K. Antidiabetic influence of dietary cumin seeds *Cuminum cyminum* in streptozotocin induced diabetic rats. Nutrition Research. 1998; 18(10):131-142.
 62. Talpur N, Echard B, Ingram C, Bagachi D, Preuss H. Effects of a novel formulation of essential oils on glucose-insulin metabolism in diabetic and hypertensive rats: a pilot study. Diabetes, Obesity and Metabolism. 2005; 7:193-199.
 63. Lee HS. Cuminaldehyde: Aldose reductase and alpha-glucosidase inhibitor derived from *Cuminum cyminum* L. seeds. Journal of Agricultural and Food Chemistry. 2005; 32:2446-2450.